arXiv:1302.0121v1 [math.RT] 1 Feb 2013

An A_∞-structure on the cohomology ring of the symmetric group S_p with coefficients in $\mathbb{F}_p$

Stephan Schmid

February 4, 2013

Abstract

Let p be a prime. Let $\mathbb{F}_p\mathrm{S}_p$ be the group algebra of the symmetric group over the finite field $\mathbb{F}_p$ with $|\mathbb{F}_p| = p$. Let $\mathbb{F}_p$ be the trivial $\mathbb{F}_p\mathrm{S}_p$-module. We present a projective resolution $\operatorname{PRes}\mathbb{F}_p$ of the module $\mathbb{F}_p$ and equip the Yoneda algebra $\operatorname{Ext}^*_{\mathbb{F}_p\mathrm{S}_p}(\mathbb{F}_p, \mathbb{F}_p)$ with an A_∞-structure such that $\operatorname{Ext}^*_{\mathbb{F}_p\mathrm{S}_p}(\mathbb{F}_p, \mathbb{F}_p)$ becomes a minimal model of the dg-algebra $\operatorname{Hom}^*_{\mathbb{F}_p\mathrm{S}_p}(\operatorname{PRes}\mathbb{F}_p, \operatorname{PRes}\mathbb{F}_p)$.

Contents

0.1 Introduction . . . 2
0.2 Outline . . . 4
0.3 Notations and conventions . . . 5

1 The projective resolution of $\mathbb{F}_p$ over $\mathbb{F}_p\mathrm{S}_p$ 7
1.1 A description of $\mathbb{Z}_{(p)}\mathrm{S}_p$ . . . 7
1.2 A projective resolution of $\mathbb{Z}_{(p)}$ over $\mathbb{Z}_{(p)}\mathrm{S}_p$ . . . 8
1.3 A projective resolution of $\mathbb{F}_p$ over $\mathbb{F}_p\mathrm{S}_p$ . . . 10

2 A_∞-algebras 11
2.1 Definitions, General theory . . . 11
2.2 The homology of $\operatorname{Hom}^*_{\mathbb{F}_p\mathrm{S}_p}(\operatorname{PRes}\mathbb{F}_p, \operatorname{PRes}\mathbb{F}_p)$ . . . 14
2.3 An A_∞-structure on $\operatorname{Ext}^*_{\mathbb{F}_p\mathrm{S}_p}(\mathbb{F}_p, \mathbb{F}_p)$ as a minimal model of $\operatorname{Hom}^*_{\mathbb{F}_p\mathrm{S}_p}$ $(\operatorname{PRes}\mathbb{F}_p, \operatorname{PRes}\mathbb{F}_p)$ . . . 18
2.4 At the prime 2 . . . 29

MSC 2010: 18G15.

0.1 Introduction

A_∞-algebras Let R be a commutative ring. Let A be a $\mathbb{Z}$-graded R-module. Let $m_1 : A \to A$ be a graded map of degree 1 with $m_1^2 = 0$, i.e. a differential on A. Let $m_2 : A \otimes A \to A$ be a graded map of degree 0 satisfying the Leibniz rule, i.e.

$$m_1 \circ m_2 = m_2 \circ (m_1 \otimes 1 + 1 \otimes m_1).$$

The map m_2 is in general not required to be associative. Instead, we require that for a morphism $m_3 : A^{\otimes 3} \to A$, the following identity holds.

$$m_2 \circ (m_2 \otimes 1 - 1 \otimes m_2) = m_1 \circ m_3 + m_3 \circ (m_1 \otimes 1^{\otimes 2} + 1 \otimes m_1 \otimes 1 + 1^{\otimes 2} \otimes m_1)$$

Following Stasheff, cf. [20], this can be continued in a certain way with higher multiplication maps to obtain a tuple of graded maps $(m_n : A^{\otimes n} \to A)_{n \geq 1}$ of certain degrees satisfying the Stasheff identities, cf. e.g. (5). The tuple $(A, (m_n)_{n \geq 1})$ is then called an A_∞-algebra.

A morphism of A_∞-algebras from $(A', (m'_n)_{n \geq 1})$ to $(A, (m_n)_{n \geq 1})$ is a tuple of graded maps $(f_n : A'^{\otimes n} \to A)_{n \geq 1}$ of certain degrees satisfying the identities (6). The first two of these are

$$\begin{aligned} &(6)[1]: & f_1 \circ m'_1 &= m_1 \circ f_1 \\ &(6)[2]: & f_1 \circ m'_2 - f_2 \circ (m'_1 \otimes 1 + 1 \otimes m'_1) &= m_1 \circ f_2 + m_2 \circ (f_1 \otimes f_1). \end{aligned}$$

So a morphism $f = (f_n)_{n \geq 1}$ of A_∞-algebras from $(A', (m'_n)_{n \geq 1})$ to $(A, (m_n)_{n \geq 1})$ contains a morphism of complexes $f_1 : (A', m'_1) \to (A, m_1)$. We say that f is a quasi-isomorphism of A_∞-algebras if f_1 is a quasi-isomorphism. Furthermore, there is a concept of homotopy for A_∞-morphisms, cf. e.g. [10, 3.7] and [14, Définition 1.2.1.7].

History The history of A_∞-algebras is outlined in [10] and [11].

As already mentioned, Stasheff introduced A_∞-algebras in 1963.

If R is a field, $\mathbb{F} := R$, we have the following basic results on A_∞-algebras, which are known since the early 1980s.

- Each quasi-isomorphism of A_∞-algebras is a homotopy equivalence, cf. [18], [9], ...
- The minimality theorem: Each A_∞-algebra $(A, (m_n)_{n \geq 1})$ is quasi-isomorphic to an A_∞-algebra $(A', \{m'_n\}_{n \geq 1})$ with $m'_1 = 0$, cf. [8], [7], [18], [4], [6], [16], ... The A_∞-algebra A' is then called a minimal model of A.

Suppose given an $\mathbb{F}$-algebra B and suppose given an B-module M together with a projective resolution $\operatorname{PRes} M$ of M. The homology of the dg-algebra $\operatorname{Hom}_B^*(\operatorname{PRes} M, \operatorname{PRes} M)$ is the Yoneda algebra $\operatorname{Ext}_B^*(M, M)$. By the minimality theorem, it is possible to construct an A_∞-structure on $\operatorname{Ext}_B^*(M, M)$ such that $\operatorname{Ext}_B^*(M, M)$ becomes a minimal model of the dg-algebra $\operatorname{Hom}_B^*(\operatorname{PRes} M, \operatorname{PRes} M)$. For the purpose of this introduction, we will call such an A_∞-structure on $\mathrm{Ext}^*_B(M,M)$ the canonical A_∞-structure on $\mathrm{Ext}^*_B(M,M)$, which is unique up to isomorphisms of A_∞-algebras, cf. [10, 3.3].

This structure has been calculated or partially calculated in several cases.

Let p be a prime.

For an arbitrary field $\mathbb{F}$, Madsen computed the canonical A_∞-structure on $\mathrm{Ext}^*_{\mathbb{F}[\alpha]/(\alpha^n)}(\mathbb{F},\mathbb{F})$, where $\mathbb{F}$ is the trivial $\mathbb{F}[\alpha]/(\alpha^n)$-module, cf. [15, Appendix B.2]. This can be used to compute the canonical A_∞-structure on the group cohomology $\mathrm{Ext}^*_{\mathbb{F}_p\mathrm{C}_m}(\mathbb{F}_p,\mathbb{F}_p)$, where $m \in \mathbb{Z}_{\geq 1}$ and C_m is the cyclic group of order m, cf. [21, Theorem 4.3.8].

Vejdemo-Johansson developed algorithms for the computation of minimal models [21]. He applied these algorithms to compute large enough parts of the canonical A_∞-structures of the group cohomologies $\mathrm{Ext}^*_{\mathbb{F}_2\mathrm{D}_8}(\mathbb{F}_2,\mathbb{F}_2)$ and $\mathrm{Ext}^*_{\mathbb{F}_2\mathrm{D}_{16}}(\mathbb{F}_2,\mathbb{F}_2)$ to distinguish them, where D_8 and D_{16} denote dihedral groups. He stated a conjecture on the complete A_∞-structure on $\mathrm{Ext}^*_{\mathbb{F}_2\mathrm{D}_8}(\mathbb{F}_2,\mathbb{F}_2)$. Furthermore, he computed parts of the canonical A_∞-structure on $\mathrm{Ext}^*_{\mathbb{F}_2\mathrm{Q}_8}(\mathbb{F}_2,\mathbb{F}_2)$ for the quaternion group Q_8. He conjecturally stated the minimal complexity of such a structure. Based on this work, there are now built-in algorithms for the Magma computer algebra system. These are capable of computing partial A_∞-structures on the group cohomology of p-groups.

In [22], Vejdemo-Johansson examined the canonical A_∞-structure $(m_n)_{n\geq 1}$ on the group cohomology $\mathrm{Ext}^*_{\mathbb{F}_p(\mathrm{C}_k\times\mathrm{C}_l)}(\mathbb{F}_p,\mathbb{F}_p)$ of the abelian group $\mathrm{C}_k \times \mathrm{C}_l$ for $k,l \geq 4$ such that k,l are multiples of p. He showed that for infinitely many $n \in \mathbb{Z}_{\geq 1}$, the operation m_n is non-zero.

In [12], Klamt investigated canonical A_∞-structures in the context of the representation theory of Lie-algebras. In particular, given certain direct sums M of parabolic Verma modules, she examined the canonical A_∞-structure $(m'_k)_{k\geq 1}$ on $\mathrm{Ext}^*_{\mathcal{O}^\mathfrak{p}}(M,M)$. She proved upper bounds for the maximal $k \in \mathbb{Z}_{\geq 1}$ such that m'_k is non-vanishing and computed the complete A_∞-structure in certain cases.

The result For $n \in \mathbb{Z}_{\geq 1}$, we denote by S_n the symmetric group on n elements.

The group cohomology $\mathrm{Ext}^*_{\mathbb{F}_p\mathrm{S}_p}(\mathbb{F}_p,\mathbb{F}_p)$ is well-known. For example, in [1, p. 74], it is calculated using group cohomological methods.

Here, we will construct the canonical A_∞-structure on $\mathrm{Ext}^*_{\mathbb{F}_p\mathrm{S}_p}(\mathbb{F}_p,\mathbb{F}_p)$.

We obtain homogeneous elements $\iota, \chi \in \mathrm{Hom}^*_{\mathbb{F}_p\mathrm{S}_p}(\mathrm{PRes}\,\mathbb{F}_p, \mathrm{PRes}\,\mathbb{F}_p) =: A$ of degree $|\iota| = 2(p-1) =: l$ and $|\chi| = l-1$ such that $\iota^j, \chi \circ \iota^j =: \chi\iota^j$ are cycles for all $j \in \mathbb{Z}_{\geq 0}$ and such that their set of homology classes $\{\overline{\iota^j} \mid j \in \mathbb{Z}_{\geq 0}\} \sqcup \{\overline{\chi\iota^j} \mid j \in \mathbb{Z}_{\geq 0}\}$ is an $\mathbb{F}_p$-basis of $\mathrm{Ext}^*_{\mathbb{F}_p\mathrm{S}_p}(\mathbb{F}_p,\mathbb{F}_p) = \mathrm{H}^*A$, cf. Proposition 20.

For all primes p, the canonical A_∞-structure $(m'_n : (\mathrm{H}^*A)^{\otimes n} \to \mathrm{H}^*A)_{n\geq 1}$ on H^*A is given as follows.

On the elements $\overline{\chi^{a_1}\iota^{j_1}} \otimes \cdots \otimes \overline{\chi^{a_n}\iota^{j_n}}$, $n \in \mathbb{Z}_{\geq 1}$, $a_i \in \{0,1\}$ and $j_i \in \mathbb{Z}_{\geq 0}$ for $i \in \{1,\ldots,n\}$, the maps m'_n are given as follows, cf. Definition 23 and Remark 37.

If there is an $i \in \{1, \ldots, n\}$ such that $a_i = 0$, then

$$\begin{aligned} m'_n(\overline{\chi^{a_1}\iota^{j_1}} \otimes \cdots \otimes \overline{\chi^{a_n}\iota^{j_n}}) &= 0 \qquad\qquad \text{for } n \neq 2 \text{ and} \\ m'_2(\overline{\chi^{a_1}\iota^{j_1}} \otimes \overline{\chi^{a_2}\iota^{j_2}}) &= \overline{\chi^{a_1+a_1}\iota^{j_1+j_2}}. \end{aligned}$$

If all a_i equal 1, then

$$\begin{aligned} m'_n(\overline{\chi\iota^{j_1}} \otimes \cdots \otimes \overline{\chi\iota^{j_n}}) &= 0 \qquad\qquad \text{for } n \neq p \text{ and} \\ m'_p(\overline{\chi\iota^{j_1}} \otimes \cdots \otimes \overline{\chi\iota^{j_p}}) &= (-1)^p\overline{\iota^{p-1+j_1+\ldots+j_p}}. \end{aligned}$$

In particular, we have $m'_n = 0$ for all $n \in \mathbb{Z}_{\geq 1} \setminus \{2, p\}$.

0.2 Outline

Section 1 The goal of section 1 is to obtain a projective resolution of the trivial $\mathbb{F}_p\mathrm{S}_p$-Specht module $\mathbb{F}_p$. A well-known method for that is "Walking around the Brauer tree", cf. [3]. Instead, we use locally integral methods to obtain a projective resolution in an explicit and straightforward manner.

Over $\mathbb{Q}$, the Specht modules are absolutely simple. Therefore we have a morphism of $\mathbb{Z}_{(p)}$-algebras $r : \mathbb{Z}_{(p)}\mathrm{S}_p \to \prod_{\lambda \dashv p} \mathrm{End}_{\mathbb{Z}_{(p)}} S^{\lambda}_{\mathbb{Z}_{(p)}} =: \Gamma$ induced by the operation of the elements of $\mathbb{Z}_{(p)}\mathrm{S}_p$ on the Specht modules S^{λ} for partitions λ of p, which becomes an Wedderburn isomorphism when tensoring with $\mathbb{Q}$. So Γ is a product of matrix rings over $\mathbb{Z}_{(p)}$. There is a well-known description of $\mathrm{im}\, r =: \Lambda$, which we use for $p \geq 3$ to obtain projective Λ-modules $\tilde{P}_k \subseteq \Lambda$, $k \in [1, p-1]$, and to construct the indecomposible projective resolution $\mathrm{PRes}\,\mathbb{Z}_{(p)}$ of the trivial $\mathbb{Z}_{(p)}\mathrm{S}_p$-Specht module $\mathbb{Z}_{(p)}$. The non-zero parts of $\mathrm{PRes}\,\mathbb{Z}_{(p)}$ are periodic with period length $l = 2(p-1)$. In section 1.3, we reduce $\mathrm{PRes}\,\mathbb{Z}_{(p)}$ modulo p to obtain a projective resolution $\mathrm{PRes}\,\mathbb{F}_p$ of the trivial $\mathbb{F}_p\mathrm{S}_p$-Specht module $\mathbb{F}_p$.

Section 2 The goal of section 2 is to compute a minimal model of the dg-algebra $\mathrm{Hom}^*_{\mathbb{F}_p\mathrm{S}_p}(\mathrm{PRes}\,\mathbb{F}_p, \mathrm{PRes}\,\mathbb{F}_p) =: A$ by equipping its homology $\mathrm{Ext}^*_{\mathbb{F}_p\mathrm{S}_p}(\mathbb{F}_p, \mathbb{F}_p) = \mathrm{H}^*A$ with a suitable A_∞-structure and finding a quasi-isomorphism of A_∞-algebras from H^*A to A.

Towards that end, we recall the basic definitions concerning A_∞-algebras and some general results in section 2.1.

While there does not seem to be a substantial difference between the cases $p = 2$ and $p \geq 3$, we separate them to simplify notation and argumentation. Consider the case $p \geq 3$. In section 2.2, we obtain a set of cycles $\{\iota^j \mid j \in \mathbb{Z}_{\geq 0}\} \cup \{\chi\iota^j \mid j \in \mathbb{Z}_{\geq 0}\}$ in A such that their homology classes are a graded basis of H^*A. In section 2.3, we obtain a suitable A_∞-structure on H^*A and a quasi-isomorphism of A_∞-algebras from H^*A to A. For the prime 2, both steps are combined in the short section 2.4.

0.3 Notations and conventions

Stipulations

- **For the remainder of this document, p will be a prime with $p \geq 3$.**
- **Write $l := 2(p-1)$.** This will give the period length of the constructed projective resolution of $\mathbb{F}_p$ over $\mathbb{F}_p\mathrm{S}_p$, cf. e.g. (1), Lemma 6.

Miscellaneous

- Concerning "∞", we assume the set $\mathbb{Z} \cup \{\infty\}$ to be ordered in such a way that ∞ is greater than any integer, i.e. $\infty > z$ for all $z \in \mathbb{Z}$, and that the integers are ordered as usual.
- For $a \in \mathbb{Z}$, $b \in \mathbb{Z} \cup \{\infty\}$, we denote by $[a, b] := \{z \in \mathbb{Z} \mid a \leq z \leq b\} \subseteq \mathbb{Z}$ the integral interval. In particular, we have $[a, \infty] = \{z \in \mathbb{Z} \mid z \geq a\} \subseteq \mathbb{Z}$ for $a \in \mathbb{Z}$.
- For $n \in \mathbb{Z}_{\geq 0}$, $k \in \mathbb{Z}$, let the binomial coefficient $\binom{n}{k}$ be defined by the number of subsets of the set $\{1, \ldots, n\}$ that have cardinality k. In particular, if $k < 0$ or $k > n$, we have $\binom{n}{k} = 0$. Then the formula $\binom{n}{k-1} + \binom{n}{k} = \binom{n+1}{k}$ holds for all $k \in \mathbb{Z}$.
- For a commutative ring R, an R-module M and $a, b \in M$, $c \in R$, we write
$$b \equiv_c a \qquad :\Longleftrightarrow \qquad a - b \in cM.$$
Often we have $M = R$ as module over itself.
- Modules are right-modules unless otherwise specified.
- For sets, we denote by $\sqcup$ the disjoint union of sets.
- $|\cdot|$: For a homogeneous element x of a graded module or a graded map g between graded modules, we denote by $|x|$ resp. $|g|$ their degrees (This is not unique for $x = 0$ resp. $g = 0$). For y a real number, $|y|$ denotes its absolute value.

Symmetric Groups Let $n \in \mathbb{Z}_{\geq 1}$. We denote the symmetric group von n elements by S_n. For a partition $\lambda \dashv n$, we denote the corresponding Specht module by S^λ.

Complexes Let R be a commutative ring and B an R-algebra.

- For a complex of B-modules
$$\cdots \to C_{k+1} \xrightarrow{d_{k+1}} C_k \xrightarrow{d_k} C_{k-1} \to \cdots \ ,$$
its k-th boundaries, cycles and homology groups are defined by $\mathrm{B}^k := \operatorname{im} d_{k+1}$, $\mathrm{Z}^k := \ker d_k$ and $\mathrm{H}^k := \mathrm{Z}^k / \mathrm{B}^k$.

  For a cycle $x \in \mathrm{Z}^k$, we denote by $\overline{x} := x + \mathrm{B}^k \in \mathrm{H}^k$ its equivalence class in homology.

- Let

$$C = (\cdots \to C_{k+1} \xrightarrow{d_{k+1}} C_k \xrightarrow{d_k} C_{k-1} \to \cdots)$$
$$C' = (\cdots \to C'_{k+1} \xrightarrow{d'_{k+1}} C'_k \xrightarrow{d'_k} C'_{k-1} \to \cdots)$$

be two complexes of B-modules.

Given $z \in \mathbb{Z}$, let

$$\mathrm{Hom}_B^z(C, C') := \prod_{i \in \mathbb{Z}} \mathrm{Hom}_B(C_{i+z}, C'_i).$$

For an additional complex $C'' = (\cdots \to C''_{k+1} \xrightarrow{d''_{k+1}} C''_k \xrightarrow{d''_k} C''_{k-1} \to \cdots)$ and maps $h = (h_i)_{i\in\mathbb{Z}} \in \mathrm{Hom}_B^m(C, C')$, $h' = (h'_i)_{i\in\mathbb{Z}} \in \mathrm{Hom}_B^n(C', C'')$, $m, n \in \mathbb{Z}$, we define the composition by component-wise composition as

$$h' \circ h := (h'_i \circ h_{i+n})_{i\in\mathbb{Z}} \in \mathrm{Hom}_B^{m+n}(C, C'').$$

We will assemble elements of $\mathrm{Hom}_B^z(C, C')$ as sums of their non-zero components, which motivates the following notations regarding "extensions by zero" and sums.

For a map $g : C_x \to C'_y$, we define $\lfloor g \rfloor_x^y \in \mathrm{Hom}_B^{x-y}(C, C')$ by

$$(\lfloor g \rfloor_x^y)_i := \begin{cases} g & \text{for } i = y \\ 0 & \text{for } i \in \mathbb{Z} \setminus \{y\} \end{cases}.$$

Let $k \in \mathbb{Z}$. Let I be a (possibly infinite) set. Let $g_i = (g_{i,j})_j \in \mathrm{Hom}_B^k(C, C')$ for $i \in I$ such that $\{i \in I \mid g_{i,j} \neq 0\}$ is finite for all $j \in \mathbb{Z}$.
We define the sum $\sum_{i\in I} g_i \in \mathrm{Hom}_B^k(C, C')$ by

$$\left(\sum\nolimits_{i\in I} g_i\right)_j := \sum_{i\in I, g_{i,j}\neq 0} g_{i,j}\,.$$

The graded R-module $\mathrm{Hom}_B^*(C, C') := \bigoplus_{k\in\mathbb{Z}} \mathrm{Hom}_B^k(C, C')$ becomes a complex via the differential $d_{\mathrm{Hom}_B^*(C,C')}$, which is defined on elements $g \in \mathrm{Hom}_B^k(C, C')$, $k \in \mathbb{Z}$ by

$$d_{\mathrm{Hom}_B^*(C,C')}(g) := d' \circ g - (-1)^k g \circ d \in \mathrm{Hom}_B^{k+1}(C, C'),$$

where $d := (d_{i+1})_{i\in\mathbb{Z}} = \sum_{i\in\mathbb{Z}} \lfloor d_{i+1} \rfloor_{i+1}^i \in \mathrm{Hom}_B^1(C, C)$ and analogously $d' := (d'_{i+1})_{i\in\mathbb{Z}} = \sum_{i\in\mathbb{Z}} \lfloor d'_{i+1} \rfloor_{i+1}^i \in \mathrm{Hom}_B^1(C', C')$.

An element $h \in \mathrm{Hom}_B^0(C, C')$ is called a complex morphism if it satisfies $d_{\mathrm{Hom}_B^*(C,C')}(h) = 0$, i.e. $d' \circ g = g \circ d$.

1 The projective resolution of $\mathbb{F}_p$ over $\mathbb{F}_p\mathrm{S}_p$

1.1 A description of $\mathbb{Z}_{(p)}\mathrm{S}_p$

Recall that $p \geq 3$ is a prime.

For R a ring and $\lambda \dashv p$ a partition of p, the $R\mathrm{S}_p$-Specht module S^λ is finitely generatey free over R with dimension independent of R, cf. [5, 8.1, proof of 8.4]. We denote this dimension by n_λ.

A partition of the form $\lambda^k := (p-k+1, 1^{k-1})$, $k \in [1,p]$ is called a *hook partition* of p.

Over the valuation ring $\mathbb{Z}_{(p)}$, there is a well-known description of the group algebra $\mathbb{Z}_{(p)}\mathrm{S}_p$, cf. e.g. [13, Corollary 4.2.8] (using [17]), cf. also [19, Chapter 7]:

Proposition 1. *Set* $n^k_{\mathrm{b}} = \binom{p-2}{k-1}$, $n^k_{\mathrm{c}} = \binom{p-2}{k-2}$. *Then* $n^k_{\mathrm{b}} + n^k_{\mathrm{c}} = \binom{p-1}{k-1} = n_{\lambda^k}$. *Set* $\Gamma := \prod_{\lambda \dashv p} \mathbb{Z}_{(p)}^{n_\lambda \times n_\lambda}$. *For* $\rho \in \Gamma$, *and* $\lambda \dashv p$, *we denote by* ρ^λ *the* λ*-th component of* ρ. *For* $\lambda = \lambda^k$, $k \in [1,p]$, *a hook partition, we name certain subblocks of* ρ^{λ^k} *as follows.*

$$\rho^{\lambda^k} = \begin{array}{cl} \begin{array}{cc} \overbrace{\qquad}^{n^k_{\mathrm{c}},} & \overbrace{\qquad}^{n^k_{\mathrm{b}}\ \textit{columns}} \end{array} & \\ \left(\begin{array}{c|c} \rho^{\lambda^k}_{\mathrm{cc}} & \rho^{\lambda^k}_{\mathrm{bc}} \\ \hline \rho^{\lambda^k}_{\mathrm{cb}} & \rho^{\lambda^k}_{\mathrm{bb}} \end{array}\right) & \begin{array}{l} \big\}\, n^k_{\mathrm{c}}\ \textit{rows} \\ \big\}\, n^k_{\mathrm{b}}\ \textit{rows} \end{array} \end{array} .$$

We have the following $\mathbb{Z}_{(p)}$*-subalgebra* Λ *of* Γ.

$$\Lambda := \{\rho \in \Gamma \mid \rho^{\lambda^k}_{\mathrm{bb}} \equiv_p \rho^{\lambda^{k+1}}_{\mathrm{cc}} \textit{ for } k \in [1,p-1] \textit{ and } \rho^{\lambda^k}_{\mathrm{bc}} \equiv_p 0 \textit{ for } k \in [1,p]\}$$

Now there is an isomorphism of $\mathbb{Z}_{(p)}$*-algebras*

$$r:\ \mathbb{Z}_{(p)}\mathrm{S}_p \xrightarrow{\sim} \Lambda.$$

such that $\rho \in \Lambda$ *acts on the trivial* $\mathbb{Z}_{(p)}\mathrm{S}_p$*-module* $\mathbb{Z}_{(p)}$ *by multiplication with its (*1×1 */ scalar-)component* ρ^{λ^1}, *i.e. for* $x \in \mathbb{Z}_{(p)}\mathrm{S}_p$ *and for* $y \in \mathbb{Z}_{(p)}$, *we have* $yx = y \cdot r(x)^{\lambda^1}$.

Example 2. For $p = 5$, the $\mathbb{Z}_{(p)}$-algebra $\mathbb{Z}_{(5)}\,\mathrm{S}_5$ is isomorphic to the subalgebra Λ of $\Gamma = \mathbb{Z}_{(5)}^{1\times 1} \times \mathbb{Z}_{(5)}^{4\times 4} \times \mathbb{Z}_{(5)}^{6\times 6} \times \mathbb{Z}_{(5)}^{4\times 4} \times \mathbb{Z}_{(5)}^{1\times 1} \times \mathbb{Z}_{(5)}^{5\times 5} \times \mathbb{Z}_{(5)}^{5\times 5}$ described as



An entry in this tuple of matrices indicates that an element of Λ must have its corresponding entry in the indicated set. A relation " $\overset{5}{\rule{2em}{0.4pt}}$ " between (equal sized) subblocks indicates that these subblocks are equivalent modulo 5, i.e. the difference of corresponding entries is an element of $5\mathbb{Z}_{(5)}$. The blocks are labeled with the diagrams of the corresponding partitions. The right ideals $\tilde{P}_i = \tilde{e}_i\Lambda$, $i \in [1,4] = [1,p-1]$ (cf. the definitions below) are framed with red lines.

1.2 A projective resolution of $\mathbb{Z}_{(p)}$ over $\mathbb{Z}_{(p)}\mathrm{S}_p$

Recall that $p \geq 3$ is a prime.

Recall from Proposition 1 that Λ is a subring of $\Gamma = \prod_{\lambda \dashv p} \mathbb{Z}_{(p)}^{n_\lambda \times n_\lambda}$.

For $\lambda \dashv p$ and $i, j \in [1, n_\lambda]$, we set $\eta_{\lambda,i,j}$ to be the element of Γ such that $(\eta_{\lambda,i,j})^{\tilde{\lambda}} = 0$ for $\tilde{\lambda} \neq \lambda$ and $(\eta_{\lambda,i,j})^{\lambda} \in \mathbb{Z}^{n_\lambda \times n_\lambda}$ has entry 1 at position (i,j) and zeros elsewhere.

Let $k \in [1, p-1]$. We obtain the idempotent

$$\tilde{e}_k := \eta_{\lambda^k, n_{\mathrm{c}}^k+1, n_{\mathrm{c}}^k+1} + \eta_{\lambda^{k+1},1,1} \in \Lambda.$$

We define corresponding projective right Λ-modules

$$\tilde{P}_k := \tilde{e}_k\Lambda \quad \text{for } k \in [1, p-1].$$

Remark 3. Let A be an R-algebra and let $e, e' \in A$ be two idempotents. For the right modules eA, $e'A$, we have the isomorphism of R-Modules

$$\begin{array}{rcl} \mathrm{Hom}_A(eA, e'A) & \overset{T_{e',e}}{\underset{\sim}{\longrightarrow}} & e'Ae \\ f & \longmapsto & T_{e',e}(f) := f(e) \\ T^{-1}_{e',e}(e'be) := (ea \mapsto e'bea) & \longleftarrow & e'be \end{array} .$$

Thus given $m \in e'Ae$, the morphism $T^{-1}_{e',e}(m)$ acts on elements $x \in eA$ by the multiplication of m on the left: $(T^{-1}_{e',e}(m))(x) = m \cdot x$.

Given idempotents $e, e', e'' \in A$, and elements $f \in \mathrm{Hom}_A(eA, e'A)$, $g \in A(e'A, e''A)$, we have $T_{e'',e}(g \circ f) = g(f(e)) = g(e'f(e)) = g(e') \cdot f(e) = T_{e'',e'}(g) \cdot T_{e',e}(f)$.

Definition 4. We define via Remark 3

$$\begin{array}{llll} \hat{e}_k := T^{-1}_{\tilde{e}_k,\tilde{e}_k}(\tilde{e}_k) & \in \mathrm{Hom}_\Lambda(\tilde{P}_k, \tilde{P}_k) & \text{for } k \in [1, p-1] \\ \hat{e}_{1,1} := T^{-1}_{\tilde{e}_1,\tilde{e}_1}(p\eta_{\lambda^1,1,1}) & \in \mathrm{Hom}_\Lambda(\tilde{P}_1, \tilde{P}_1) & \\ \hat{e}_{p-1,p-1} := T^{-1}_{\tilde{e}_{p-1},\tilde{e}_{p-1}}(p\eta_{\lambda^p,1,1}) & \in \mathrm{Hom}_\Lambda(\tilde{P}_{p-1}, \tilde{P}_{p-1}) & \\ \hat{e}_{k+1,k} := T^{-1}_{\tilde{e}_{k+1},\tilde{e}_k}(\eta_{\lambda^{k+1},n_c^{k+1}+1,1}) & \in \mathrm{Hom}_\Lambda(\tilde{P}_k, \tilde{P}_{k+1}) & \text{for } k \in [1, p-2] \\ \hat{e}_{k,k+1} := T^{-1}_{\tilde{e}_k,\tilde{e}_{k+1}}(p\eta_{\lambda^{k+1},1,n_c^{k+1}+1}) & \in \mathrm{Hom}_\Lambda(\tilde{P}_{k+1}, \tilde{P}_k) & \text{for } k \in [1, p-2]. \end{array}$$

Note that $\hat{e}_k$ is the identity map on $\tilde{P}_k$ for $k \in [1, p-1]$.

Moreover, we define the $\mathbb{Z}_{(p)}\mathrm{S}_p$-linear map $\hat{\varepsilon} : \tilde{P}_1 \to \mathbb{Z}_{(p)}$, $\hat{\varepsilon}(\rho) := \rho^{\lambda^1}$.

It is straightforward to show the following lemma.

Lemma 5. *We have*

$$\begin{array}{rcll} \hat{e}_{1,1} + \hat{e}_{1,2} \circ \hat{e}_{2,1} & = & p\hat{e}_1 & \\ \hat{e}_{k,k-1} \circ \hat{e}_{k-1,k} + \hat{e}_{k,k+1} \circ \hat{e}_{k+1,k} & = & p\hat{e}_k & \textit{for } k \in [2, p-2] \\ \hat{e}_{p-1,p-2} \circ \hat{e}_{p-2,p-1} + \hat{e}_{p-1,p-1} & = & p\hat{e}_{p-1} & \\ \hat{\varepsilon} \circ \hat{e}_{1,1} & = & p\hat{\varepsilon}. & \end{array}$$

Furthermore, it is straightforward to check that we obtain a projective resolution of $\mathbb{Z}_{(p)}$ as follows. We set

$$\tilde{\mathrm{Pr}}_i := \begin{cases} \tilde{P}_{\omega(i)} & i \geq 0 \\ 0 & i < 0 \end{cases},$$

where the integer $\omega(i)$ is given by the following construction: Recall the stipulation $l := 2(p-1)$. We have $i = jl + r$ for some $j \in \mathbb{Z}$ and $0 \leq r \leq l-1$. Then

$$\omega(i) := \begin{cases} r+1 & \text{for } 0 \leq r \leq p-2 \\ l - r = 2(p-1) - r & \text{for } p-1 \leq r \leq 2(p-1) - 1 = l-1 \end{cases}. \tag{1}$$

So $\omega(i)$ increases by steps of one from 1 to $p-1$ as i runs from jl to $jl+(p-2)$ and $\omega(i)$ decreases from $p-1$ to 1 as i runs from $jl+(p-1)$ to $jl+(l-1)$. Finally we set

$$\hat{d}_i := \begin{cases} \hat{e}_{\omega(i-1),\omega(i)} : \tilde{P}_{\omega(i)} \to \tilde{P}_{\omega(i-1)} & i \geq 1 \\ 0 & i \leq 0 \end{cases}.$$

Now we have the projective resolution of $\mathbb{Z}_{(p)}$

$$\cdots \xrightarrow{\hat{d}_3} \tilde{\mathrm{Pr}}_2 \xrightarrow{\hat{d}_2} \tilde{\mathrm{Pr}}_1 \xrightarrow{\hat{d}_1} \tilde{\mathrm{Pr}}_0 \xrightarrow{0=\hat{d}_0} 0 \to \cdots , \tag{2}$$

written more explicitly as

$$\begin{aligned} \cdots \to \tilde{P}_2 \xrightarrow{\hat{e}_{1,2}} \tilde{P}_1 \xrightarrow{\hat{e}_{1,1}} \tilde{P}_1 \xrightarrow{\hat{e}_{2,1}} \tilde{P}_2 \to \ldots \to \tilde{P}_{p-2} \xrightarrow{\hat{e}_{p-1,p-2}} \tilde{P}_{p-1} \\ \xrightarrow{\hat{e}_{p-1,p-1}} \tilde{P}_{p-1} \xrightarrow{\hat{e}_{p-2,p-1}} \tilde{P}_{p-2} \to \ldots \to \tilde{P}_2 \xrightarrow{\hat{e}_{1,2}} \tilde{P}_1 \to 0 \to \cdots , \end{aligned}$$

with augmentation $\hat{\varepsilon} : \tilde{P}_1 \to \mathbb{Z}_{(p)}$.

1.3 A projective resolution of $\mathbb{F}_p$ over $\mathbb{F}_p\mathrm{S}_p$

The isomorphism $r : \mathbb{Z}_{(p)}\mathrm{S}_p \to \Lambda$ from Proposition 1 induces an isomorphism of $\mathbb{F}_p$-algebras $\mathbb{F}_p\mathrm{S}_p = \mathbb{Z}_{(p)}\mathrm{S}_p/(p\mathbb{Z}_{(p)}\mathrm{S}_p) \xrightarrow{\bar{r}} \Lambda/(p\Lambda) =: \bar{\Lambda}$.
For the sake of simplicity in the next step, we identify $\bar{\Lambda}$ and $\mathbb{F}_p\mathrm{S}_p$ along $\bar{r}$.

Lemma 6. *Recall that* $p \geq 3$ *is a prime. Applying the functor* $- \underset{\Lambda}{\otimes} {}_{\Lambda}\bar{\Lambda}_{\bar{\Lambda}}$ *, we obtain*

- *the projective modules* $P_k := \tilde{P}_k \underset{\Lambda}{\otimes} {}_{\Lambda}\bar{\Lambda}_{\bar{\Lambda}}$ *for* $k \in [1, p-1]$,
- $\mathbb{F}_p := \mathbb{Z}_{(p)} \underset{\Lambda}{\otimes} {}_{\Lambda}\bar{\Lambda}_{\bar{\Lambda}}$ *(the trivial* $\mathbb{F}_p\mathrm{S}_p$*-module),*
- $$\begin{array}{llll} e_k & := \hat{e}_k \underset{\Lambda}{\otimes} {}_{\Lambda}\bar{\Lambda}_{\bar{\Lambda}} & \in \mathrm{Hom}_{\mathbb{F}_p\mathrm{S}_p}(P_k, P_k) & \text{for } k \in [1, p-1], \\ e_{1,1} & := \hat{e}_{1,1} \underset{\Lambda}{\otimes} {}_{\Lambda}\bar{\Lambda}_{\bar{\Lambda}} & \in \mathrm{Hom}_{\mathbb{F}_p\mathrm{S}_p}(P_1, P_1), & \\ e_{p-1,p-1} & := \hat{e}_{p-1,p-1} \underset{\Lambda}{\otimes} {}_{\Lambda}\bar{\Lambda}_{\bar{\Lambda}} & \in \mathrm{Hom}_{\mathbb{F}_p\mathrm{S}_p}(P_{p-1}, P_{p-1}), & \\ e_{k+1,k} & := \hat{e}_{k+1,k} \underset{\Lambda}{\otimes} {}_{\Lambda}\bar{\Lambda}_{\bar{\Lambda}} & \in \mathrm{Hom}_{\mathbb{F}_p\mathrm{S}_p}(P_k, P_{k+1}) & \text{for } k \in [1, p-2], \\ e_{k,k+1} & := \hat{e}_{k,k+1} \underset{\Lambda}{\otimes} {}_{\Lambda}\bar{\Lambda}_{\bar{\Lambda}} & \in \mathrm{Hom}_{\mathbb{F}_p\mathrm{S}_p}(P_{k+1}, P_k) & \text{for } k \in [1, p-2], \\ \varepsilon & := \hat{\varepsilon} \underset{\Lambda}{\otimes} {}_{\Lambda}\bar{\Lambda}_{\bar{\Lambda}} & \in \mathrm{Hom}_{\mathbb{F}_p\mathrm{S}_p}(P_1, \mathbb{F}_p). & \end{array}$$

The complex

$$\mathrm{PRes}\,\mathbb{F}_p := (\mathrm{PRes}\,\mathbb{Z}_{(p)}) \underset{\Lambda}{\otimes} {}_{\Lambda}\bar{\Lambda}_{\bar{\Lambda}} = (\cdots \xrightarrow{d_3} \mathrm{Pr}_2 \xrightarrow{d_2} \mathrm{Pr}_1 \xrightarrow{d_1} \mathrm{Pr}_0 \xrightarrow{0=d_0} 0 \to \cdots), \tag{3}$$

$$\mathrm{Pr}_i := \begin{cases} P_{\omega(i)} & i \geq 0 \\ 0 & i < 0 \end{cases} \qquad d_i := \begin{cases} e_{\omega(i-1),\omega(i)} : P_{\omega(i)} \to P_{\omega(i-1)} & i \geq 1 \\ 0 & i \leq 0, \end{cases}$$

is a projective resolution of $\mathbb{F}_p$ *with augmentation* $\varepsilon : P_1 \to \mathbb{F}_p$. *More explicitly,* $\operatorname{PRes}\mathbb{F}_p$ *is*

$$\begin{aligned} \ldots \to \underbrace{P_2}_{l+1} \xrightarrow{e_{1,2}} \underbrace{P_1}_{l=2(p-1)} \xrightarrow{e_{1,1}} \underbrace{P_1}_{(p-2)+p-1} \xrightarrow{e_{2,1}} \underbrace{P_2}_{(p-2)+p-2} \to \ldots \to \underbrace{P_{p-2}}_{p=(p-2)+2} \xrightarrow{e_{p-1,p-2}} \underbrace{P_{p-1}}_{(p-2)+1} \\ \xrightarrow{e_{p-1,p-1}} \underbrace{P_{p-1}}_{p-2} \xrightarrow{e_{p-2,p-1}} \underbrace{P_{p-2}}_{p-3} \to \ldots \to \underbrace{P_2}_{1} \xrightarrow{e_{1,2}} \underbrace{P_1}_{0} \to 0. \end{aligned}$$

Lemma 7. *Recall that* $p \geq 3$ *is a prime.*

(a) *We have the relations*

$$\begin{array}{lcll} e_{1,1} + e_{1,2} \circ e_{2,1} & = & 0 & \\ e_{k,k-1} \circ e_{k-1,k} + e_{k,k+1} \circ e_{k+1,k} & = & 0 & \text{for } k \in [2, p-2] \\ e_{p-1,p-2} \circ e_{p-2,p-1} + e_{p-1,p-1} & = & 0 & \\ \varepsilon \circ e_{1,1} & = & 0 & \end{array}$$

and e_k *is the identity on* P_k *for* $k \in [1, p-1]$.

(b) *Given* $k \in [2, p-1]$, *we have* $\operatorname{Hom}_{\mathbb{F}_p \mathrm{S}_p}(P_k, \mathbb{F}_p) = \{0\}$.

(c) *Given* $k, k' \in [1, p-1]$ *such that* $|k - k'| > 1$, *we have* $\operatorname{Hom}_{\mathbb{F}_p \mathrm{S}_p}(P_k, P_{k'}) = \{0\}$.

(d) *The set* $\{\varepsilon\}$ *is an* $\mathbb{F}_p$*-basis of* $\operatorname{Hom}_{\mathbb{F}_p \mathrm{S}_p}(P_1, \mathbb{F}_p)$.

Assertion (a) results from Lemma 5.

Assertions (b), (c) and (d) are derived from corresponding assertions over $\mathbb{Z}_{(p)}\mathrm{S}_p$ using $\operatorname{Hom}_{\mathbb{F}_p \mathrm{S}_p}(P/pP, M/pM) \simeq \operatorname{Hom}_{\mathbb{Z}_{(p)}\mathrm{S}_p}(P, M)/p \operatorname{Hom}_{\mathbb{Z}_{(p)}\mathrm{S}_p}(P, M)$ for $\mathbb{Z}_{(p)}\mathrm{S}_p$-modules P and M, where P is projective.

2 A_∞-algebras

2.1 Definitions, General theory

In this subsection, we review results presented in [10] and we fix notation.

Let R be a commutative ring. We understand linear maps between R-modules to be R-linear. Tensor products are tensor products over R. By graded R-modules we understand $\mathbb{Z}$-graded R-modules.

Definition 8. In the definition of the tensor product of graded maps, we implement the *Koszul sign rule*: Let A_1, A_2, B_1, B_2 be graded R-modules and $g : A_1 \to B_1$, $h : A_2 \to B_2$ graded maps. Then we set for homogeneous elements $x \in A_1, y \in A_2$

$$(g \otimes h)(x \otimes y) := (-1)^{|h| \cdot |x|} g(x) \otimes h(y). \tag{4}$$

Concerning the signs in the definition of A$_\infty$-algebras and A$_\infty$-morphisms, we follow the variant given e.g. in [14] and [7].

Definition 9. Let $n \in \mathbb{Z}_{\geq 0} \cup \{\infty\}$.

(i) Let A be a graded R-module. A *pre-*A$_n$*-structure on* A is a family of graded maps $(m_k : A^{\otimes k} \to A)_{k\in[1,n]}$ with $|m_k| = 2 - k$ for $k \in [1, n]$. The tuple $(A, (m_k)_{k\in[1,n]})$ is called a pre-A$_n$-algebra.

(ii) Let A, A' be graded R-modules. A *pre-*A$_n$*-morphism from* A' *to* A is a family of graded maps $(f_k : A'^{\otimes k} \to A)_{k\in[1,n]}$ with $|f_k| = 1 - k$ for $k \in [1, n]$.

Definition 10. Let $n \in \mathbb{Z}_{\geq 0} \cup \{\infty\}$.

(i) An A$_n$*-algebra* is a pre-A$_n$-algebra $(A, (m_k)_{k\in[1,n]})$ such that for $k \in [1, n]$

$$\sum_{\substack{k=r+s+t,\\ r,t\geq 0, s\geq 1}} (-1)^{rs+t} m_{r+1+t} \circ (1^{\otimes r} \otimes m_s \otimes 1^{\otimes t}) = 0. \qquad (5)[k]$$

In abuse of notation, we sometimes abbreviate $A = (A, (m_k)_{k\geq 1})$ for A$_\infty$-algebras.

(ii) Let $(A', (m'_k)_{k\in[1,n]})$ and $(A, (m_k)_{k\in[1,n]})$ be A$_n$-algebras. An A$_n$*-morphism* or *morphism of* A$_n$*-algebras* from $(A', (m'_k)_{k\in[1,n]})$ to $(A, (m_k)_{k\in[1,n]})$ is a pre-A$_n$-morphism $(f_k)_{k\in[1,n]}$ such that for $k \in [1, n]$, we have

$$\sum_{\substack{k=r+s+t\\ r,t\geq 0, s\geq 1}} (-1)^{rs+t} f_{r+1+t} \circ (1^{\otimes r} \otimes m'_s \otimes 1^{\otimes t}) = \sum_{\substack{1\leq r\leq k\\ i_1+\ldots+i_r=k\\ i_s\geq 1}} (-1)^v m_r \circ (f_{i_1} \otimes f_{i_2} \otimes \ldots \otimes f_{i_r}), \qquad (6)[k]$$

where $v := \sum_{1\leq t<s\leq r} (1 - i_s) i_t$.

Example 11 (dg-algebras)**.** Let $(A, (m_k)_{k\geq 1})$ be an A$_\infty$-algebra. If $m_n = 0$ for $n \geq 3$ then A is called a *differential graded algebra* or *dg-algebra*. In this case the equations (5)[n] for $n \geq 4$ become trivial: We have $(r + 1 + t) + s = n + 1 \Rightarrow (r + 1 + t) + s \geq 5$ $\Rightarrow m_{r+1+t} = 0$ or $m_s = 0$. So all summands in (5)[n] are zero for $n \geq 4$. Here are the equations for $n \in \{1, 2, 3\}$:

$$\begin{aligned}
(5)[1]: \quad & 0 = m_1 \circ m_1 \\
(5)[2]: \quad & 0 = m_1 \circ m_2 - m_2 \circ (m_1 \otimes 1 + 1 \otimes m_1) \\
(5)[3]: \quad & 0 = m_1 \circ m_3 + m_2 \circ (1 \otimes m_2 - m_2 \otimes 1) \\
& \quad + m_3 \circ (m_1 \otimes 1^{\otimes 2} + 1 \otimes m_2 \otimes 1 + 1^{\otimes 2} \otimes m_1) \\
& \overset{m_3=0}{=} m_2 \circ (1 \otimes m_2 - m_2 \otimes 1)
\end{aligned}$$

So (5)[1] ensures that m_1 is a differential. Moreover, (5)[3] states that m_2 is an associative binary operation, since for homogeneous $x, y, z \in A$ we have $0 = m_2 \circ (1 \otimes m_2 - m_2 \otimes 1)(x \otimes y \otimes z) = m_2(x \otimes m_2(y \otimes z) - m_2(x \otimes y) \otimes z)$, where because of $|m_2| = 0$ there are no additional signs caused by the Koszul sign rule. Equation (5)[2] is the Leibniz rule.

Example 12 (A_n-morphisms induce complex morphisms)**.**
Let $n \in \mathbb{Z}_{\geq 1} \cup \{\infty\}$. Let $(A', (m'_k)_{k\in[1,n]})$ and $(A, (m_k)_{k\in[1,n]})$ be two A_n-algebras and let $(f_k)_{k\in[1,n]} : (A', (m'_k)_{k\in[1,n]}) \to (A, (m_k)_{k\in[1,n]})$ be an A_n-morphism.

By (5)[1], (A', m'_1) and (A, m_1) are complexes. Equation (6)[1] is

$$f_1 \circ m'_1 = m_1 \circ f_1.$$

Thus $f_1 : (A', m'_1) \to (A, m_1)$ is a complex morphism.

For $n \geq 2$, we have also (6)[2]:

$$f_1 \circ m'_2 - f_2 \circ (m'_1 \otimes 1 + 1 \otimes m'_1) = m_1 \circ f_2 + m_2 \circ (f_1 \otimes f_1) \tag{7}$$

Recall the conventions concerning $\mathrm{Hom}^k_B(C, C')$.

Lemma 13 (cf. e.g. [10, Section 3.3])**.** *Let B be an (ordinary) R-algebra and $M = ((M_i)_{i\in\mathbb{Z}}, (d_i)_{i\in\mathbb{Z}})$ a complex of B-modules, that is a sequence $(M_i)_{i\in\mathbb{Z}}$ of B-modules and B-linear maps $d_i : M_i \to M_{i-1}$ such that $d_{i-1} \circ d_i = 0$ for all $i \in \mathbb{Z}$. Let*

$$\begin{aligned}\mathrm{Hom}^i_B(M, M) :=& \prod_{z\in\mathbb{Z}} \mathrm{Hom}_B(M_{z+i}, M_z)\\ =& \{g = (g_z)_{z\in\mathbb{Z}} \mid g_z \in \mathrm{Hom}_B(M_{z+i}, M_z) \text{ for } z \in \mathbb{Z}\}.\end{aligned}$$

Then

$$A = \mathrm{Hom}^*_B(M, M) := \bigoplus_{i\in\mathbb{Z}} \mathrm{Hom}^i_B(M, M)$$

is a graded R-module. We have $d := (d_{z+1})_{z\in\mathbb{Z}} = \sum_{i\in\mathbb{Z}} \lfloor d_{i+1} \rfloor^i_{i+1} \in \mathrm{Hom}^1_B(M, M)$. We define $m_1 := d_{\mathrm{Hom}^*(M,M)} : A \to A$, that is for homogeneous $g \in A$ we have

$$m_1(g) = d \circ g - (-1)^{|g|} g \circ d.$$

We define $m_2 : A^{\otimes 2} \to A$ for homogeneous $g, h \in A$ to be composition, i.e.

$$m_2(g \otimes h) := g \circ h.$$

For $n \geq 3$ we set $m_n : A^{\otimes n} \to A$, $m_n = 0$. Then $(m_n)_{n\geq 1}$ is an A_∞-algebra structure on $A = \mathrm{Hom}^*_B(M^*, M^*)$. More precisely, $(A, (m_n)_{n\geq 1})$ is a dg-algebra.

Remark 14. In $\mathrm{Hom}^*(\mathrm{PRes}\,\mathbb{F}_p, \mathrm{PRes}\,\mathbb{F}_p)$ we have (cf. (3))

$$d = \sum_{i\geq 0} \lfloor e_{\omega(i),\omega(i+1)} \rfloor^i_{i+1}.$$

Definition 15 (Homology of A_∞-algebras, quasi-isomorphisms, minimality, minimal models)**.** As $m_1^2 = 0$ (cf. (5)[1]) and $|m_1| = 1$, we have the complex

$$\cdots \to A^{i-1} \xrightarrow{m_1|_{A^{i-1}}} A^i \xrightarrow{m_1|_{A^i}} A^{i+1} \to \cdots .$$

We define $\mathrm{H}^k A := \ker(m_1|_{A^k}) / \operatorname{im}(m_1|_{A^{k-1}})$ and $\mathrm{H}^* A := \bigoplus_{k\in\mathbb{Z}} \mathrm{H}^k A$, which gives the homology of A the structure of a graded R-module.

A morphism of A_∞-algebras $(f_k)_{k\geq 1} : (A', (m'_k)_{k\geq 1}) \to (A, (m_k)_{k\geq 1})$ is called a *quasi-isomorphism* if the morphism of complexes $f_1 : (A', m'_1) \to (A, m_1)$ (cf. Example 12) is a quasi-isomorphism.

An A_∞-algebra is called *minimal*, if $m_1 = 0$. If A is an A_∞-algebra and A' is a minimal A_∞-algebra quasi-isomorphic to A, then A' is called a *minimal model of A*.

The existence of minimal models is assured by the following theorem.

Theorem 16. *(minimality theorem, cf. [11] (history), [8], [7], [18], [4], [6], [16], ...) Let $(A, (m_k)_{k\geq 1})$ be an A_∞-algebra such that the homology H^*A is a projective R-module. Then there exists an A_∞-algebra structure $(m'_k)_{k\geq 1}$ on H^*A and a quasi-isomorphism of A_∞-algebras $(f_k)_{k\geq 1} : (\mathrm{H}^*A, (m'_k)_{k\geq 1}) \to (A, (m_k)_{k\geq 1})$, such that*

- *$m'_1 = 0$ and*
- *the complex morphism $f_1 : (\mathrm{H}^*A, m'_1) \to (A, m_1)$ induces the identity in homology. I.e. each element $x \in \mathrm{H}^*A$, which is a homology class of (A, m_1), is mapped by f_1 to a representing cycle.*

For constructing A_∞-structures induced by another A_∞-algebra, we have the following

Lemma 17 (cf. [7, Proof of Theorem 1])**.** *Let $n \in \mathbb{Z}_{\geq 1} \cup \{\infty\}$. Let $(A', (m'_k)_{k\in[1,n]})$ be a pre-A_n-algebra. Let $(A, (m_k)_{k\in[1,n]})$ be an A_n-algebra. Let $(f_k)_{k\in[1,n]}$ be a pre-A_n-morphism from A' to A such that (6)[k] holds for $k \in [1, n]$. Suppose f_1 to be injective. Then $(A', (m'_k)_{k\in[1,n]})$ is an A_n-algebra and $(f_k)_{k\in[1,n]}$ is a morphism of A_n-algebras from $(A', (m'_k)_{k\in[1,n]})$ to $(A, (m_k)_{k\in[1,n]})$.*

This results from the bar construction and a straightforward induction on n.

Lemma 18 ([23, Theorem 5])**.** *Let R be a commutative ring and $(A, (m_n)_{n\geq 1})$ be a dg-algebra (over R). Suppose given a graded R-module B and graded maps $f_n : B^{\otimes n} \to A$, $m'_n : B^{\otimes n} \to B$ for $n \geq 1$. Suppose given $k \geq 1$ such that we have $f_i = 0$ for $i \geq k$, we have $m'_i = 0$ for $i \geq k+1$, and (6)[n] is satisfied for $n \in [1, 2k-2]$. Then (6)[n] is satisfied for all $n \geq 1$.*

2.2 The homology of $\operatorname{Hom}^*_{\mathbb{F}_p\mathrm{S}_p}(\operatorname{PRes}\mathbb{F}_p, \operatorname{PRes}\mathbb{F}_p)$

We need a well-known result of homological algebra in a particular formulation:

Lemma 19. *Let F be a field. Let B be an F-algebra. Let M be a B-module. Let $Q = (\cdots \to Q_2 \xrightarrow{d_2} Q_1 \xrightarrow{d_1} Q_0 \to 0 \to \cdots)$ be a projective resolution of M with augmentation $\varepsilon : Q_0 \to M$. Then we have maps for $k \in \mathbb{Z}$*

$$\begin{aligned} \Psi_k : \operatorname{Hom}^k_B(Q,Q) &\to \operatorname{Hom}^k_B(Q,M) := \operatorname{Hom}_B(Q_k, M) \\ (g_i : Q_{i+k} \to Q_i)_{i\in\mathbb{Z}} &\mapsto \varepsilon \circ g_0 \end{aligned}$$

The right side is equipped with the differentials (dualization of d_k)

$$\begin{aligned} (d_k)^* : \operatorname{Hom}_B(Q_k, M) &\to \operatorname{Hom}_B(Q_{k+1}, M) \\ g &\mapsto (-1)^k g \circ d_k \end{aligned}$$

and the left side is equipped with the differential m_1 of its dg-algebra structure, cf. Lemma 13.

Then $(\Psi_k)_{k\in\mathbb{Z}}$ becomes a complex morphism from the complex $\operatorname{Hom}^*_B(Q,Q)$ to the complex $\operatorname{Hom}^*_B(Q,M)$ that induces isomorphisms $\bar{\Psi}_k$ of F-vector spaces on the homology

$$\begin{aligned} \bar{\Psi}_k : \mathrm{H}^k \operatorname{Hom}^*_B(Q,Q) &\xrightarrow{\sim} \mathrm{H}^k \operatorname{Hom}^*_B(Q,M) \\ \overline{(g_i : Q_{i+k} \to Q_i)_{i\in\mathbb{Z}}} &\mapsto \overline{\varepsilon \circ g_0} \end{aligned}$$

Lemma 19 is [2, §5 Proposition 4a)] applied to the quasi-isomorphism induced by the augmentation, cf. [2, §3 Définition 1].

Recall the notation $\lfloor x \rfloor^z_y$ for the description of elements of $\operatorname{Hom}^k_B(C, C')$.

Proposition 20. *Recall that $p \geq 3$ is a prime and $l = 2(p-1)$.*
Write $A := \operatorname{Hom}^*_{\mathbb{F}_p\mathrm{S}_p}(\operatorname{PRes}\mathbb{F}_p, \operatorname{PRes}\mathbb{F}_p)$. Let

$$\begin{aligned} \iota &:= \textstyle\sum_{i\geq 0} \lfloor e_{\omega(i)} \rfloor^{i}_{i+l} = \sum_{i\geq 0} \sum_{k=0}^{l-1} \lfloor e_{\omega(k)} \rfloor^{il+k}_{(i+1)l+k} \in A^l \\ \chi &:= \textstyle\sum_{i\geq 0} \Big(\lfloor e_1 \rfloor^{il}_{il+l-1} + \Big(\sum_{k=1}^{p-2} \lfloor e_{k+1,k} \rfloor^{il+k}_{il+l-1+k} \Big) \\ &\qquad + \lfloor e_{p-1} \rfloor^{il+(p-1)}_{il+l-1+(p-1)} + \Big(\textstyle\sum_{k=1}^{p-2} \lfloor e_{p-k-1,p-k} \rfloor^{il+(p-1)+k}_{il+l-1+(p-1)+k} \Big) \Big) \in A^{l-1}. \end{aligned}$$

(a) *For $j \geq 0$, we have $\iota^j = \sum_{i\geq 0} \lfloor e_{\omega(i)} \rfloor^{i}_{i+jl} = \sum_{i\geq 0} \sum_{k=0}^{l-1} \lfloor e_{\omega(k)} \rfloor^{il+k}_{(i+j)l+k}$.*

(b) *Suppose given $y \geq 0$. Let $h \in A^y$ be l-periodic, that is $h = \sum_{i\geq 0} \sum_{k=0}^{l-1} \lfloor h_k \rfloor^{il+k}_{il+k+y}$. Then for $j \geq 0$, we have*

$$h \circ \iota^j = \iota^j \circ h = \sum_{i\geq 0} \sum_{k=0}^{l-1} \lfloor h_k \rfloor^{il+k}_{(i+j)l+k+y} \in A^{y+jl}.$$

(c) *Suppose given $y \in \mathbb{Z}$. For $h \in A^y$ and $j \geq 0$, we have $m_1(h \circ \iota^j) = m_1(h) \circ \iota^j$.*

(d) *For $j \geq 0$, we have $m_1(\iota^j) = 0$. Thus ι^j is a cycle.*

(e) *For $j \geq 0$, we have*

$$\begin{aligned}\chi\iota^j &:= \chi \circ \iota^j = \iota^j \circ \chi \\ &= \sum\nolimits_{i \geq 0} \Big(\lfloor e_1 \rfloor^{il}_{(i+j+1)l-1} + \Big(\sum\nolimits_{k=1}^{p-2} \lfloor e_{k+1,k} \rfloor^{il+k}_{(i+j+1)l-1+k} \Big) \\ &\qquad + \lfloor e_{p-1} \rfloor^{il+(p-1)}_{(i+j+1)l-1+(p-1)} + \Big(\sum\nolimits_{k=1}^{p-2} \lfloor e_{p-k-1,p-k} \rfloor^{il+(p-1)+k}_{(i+j+1)l-1+(p-1)+k} \Big) \Big) \in A^{jl+l-1}.\end{aligned}$$

For convenience, we also define $\chi^0\iota^j := \iota^j$ and $\chi^1\iota^j := \chi\iota^j = \chi \circ \iota^j$ for $j \geq 0$.

(f) *For $j \geq 0$, we have $m_1(\chi\iota^j) = 0$. Thus $\chi\iota^j$ is a cycle.*

(g) *Suppose given $k \in \mathbb{Z}$. A $\mathbb{F}_p$-basis of $\mathrm{H}^k A$ is given by*

$$\begin{aligned} &\{\overline{\iota^j}\} \text{ if } k = jl \text{ for some } j \geq 0 \\ &\{\overline{\chi\iota^j}\} \text{ if } k = jl + l - 1 \text{ for some } j \geq 0 \\ &\emptyset \text{ else.}\end{aligned}$$

Thus the set $\mathfrak{B} := \{\overline{\iota^j} \mid j \geq 0\} \sqcup \{\overline{\chi\iota^j} \mid j \geq 0\}$ is an $\mathbb{F}_p$-basis of $\mathrm{H}^* A = \bigoplus_{z \in \mathbb{Z}} \mathrm{H}^z A$.

Proof. The element ι is well-defined since $\omega(y) = \omega(l + y)$ for $y \geq 0$.
In the definition of χ we need to check that the "$\lfloor * \rfloor^*_*$" are well defined. This is easily proven by calculating the $\omega(y)$ where y is the lower respective upper index of "$\lfloor * \rfloor^*_*$".

(a): As $\mathrm{Pr}_i = \{0\}$ for $i < 0$, the identity element of A is given by $\iota^0 = \sum_{i \geq 0} \lfloor e_{\omega(i)} \rfloor^i_i$, which agrees with the assertion in case $j = 0$. So we have proven the induction basis for induction on j. So now assume that for some $j \geq 0$ the assertion holds. Then

$$\begin{aligned}\iota^{j+1} &= \iota \circ \iota^j = \Big(\sum\nolimits_{i \geq 0} \lfloor e_{\omega(i)} \rfloor^i_{i+l} \Big) \circ \Big(\sum\nolimits_{i' \geq 0} \lfloor e_{\omega(i')} \rfloor^{i'}_{i'+jl} \Big) \\ &= \sum\nolimits_{i \geq 0} \lfloor e_{\omega(i)} \circ e_{\omega(i+l)} \rfloor^i_{i+l+jl} = \sum\nolimits_{i \geq 0} \lfloor e_{\omega(i)} \rfloor^i_{i+(j+1)l} .\end{aligned}$$

Thus the proof by induction is complete.

(b): We have

$$\begin{aligned}\iota^j \circ h &= \left(\sum_{i \geq 0} \sum_{k=0}^{l-1} \lfloor e_{\omega(il+k)} \rfloor^{il+k}_{(i+j)l+k} \right) \circ \left(\sum_{i' \geq 0} \sum_{k'=0}^{l-1} \lfloor h_{k'} \rfloor^{i'l+k'}_{i'l+k'+y} \right) \overset{\substack{i' \rightsquigarrow i+j \\ k' \rightsquigarrow k}}{=} \sum_{i \geq 0} \sum_{k=0}^{l-1} \lfloor h_k \rfloor^{il+k}_{(i+j)l+k+y} \\ h \circ \iota^j &= \left(\sum_{i \geq 0} \sum_{k=0}^{l-1} \lfloor h_k \rfloor^{il+k}_{il+k+y} \right) \circ \left(\sum_{i' \geq 0} \lfloor e_{\omega(i')} \rfloor^{i'}_{i'+jl} \right) = \sum_{i \geq 0} \sum_{k=0}^{l-1} \lfloor h_k \rfloor^{il+k}_{(i+j)l+k+y} .\end{aligned}$$

So we have proven (b).

(c): The differential d of $\mathrm{PRes}\,\mathbb{F}_p$ is l-periodic (cf. Remark 14) and thus

$$m_1(h) \circ \iota^j = (d \circ h - (-1)^y h \circ d) \circ \iota^j$$

$$\overset{(b),|\iota^j|\equiv_2 0}{=} d \circ h \circ \iota^j - (-1)^{y+|\iota^j|} h \circ \iota^j \circ d = m_1(h \circ \iota^j).$$

(d): We have $m_1(\iota^j) \overset{(c)}{=} m_1(\iota^0) \circ \iota^j = (d \circ \iota^0 - (-1)^0 \iota^0 d) \circ \iota^j = (d - d) \circ \iota^j = 0$.

(e) is implied by (b) using the fact that χ is l-periodic.

(f): Because of (c) we have $m_1(\chi\iota^j) = m_1(\chi) \circ \iota^j$. Because $|\chi| = l - 1$ is odd we have

$$\begin{aligned}
&m_1(\chi) = d \circ \chi - (-1)\chi \circ d = \chi \circ d + d \circ \chi \\
&\overset{\text{R.}14}{=} \Big(\sum\nolimits_{i\geq 0} \Big(\lfloor e_1 \rfloor^{il}_{il+l-1} + \Big(\sum\nolimits_{k=1}^{p-2} \lfloor e_{k+1,k} \rfloor^{il+k}_{il+l-1+k} \Big) + \lfloor e_{p-1} \rfloor^{il+(p-1)}_{il+l-1+(p-1)} \\
&\quad + \Big(\sum\nolimits_{k=1}^{p-2} \lfloor e_{p-k-1,p-k} \rfloor^{il+(p-1)+k}_{il+l-1+(p-1)+k} \Big) \Big) \Big) \circ \Big(\sum\nolimits_{y\geq 0} \lfloor e_{\omega(y),\omega(y+1)} \rfloor^{y}_{y+1} \Big) \\
&\quad + \Big(\sum\nolimits_{y\geq 0} \lfloor e_{\omega(y),\omega(y+1)} \rfloor^{y}_{y+1} \Big) \circ \Big(\sum\nolimits_{i\geq 0} \Big(\lfloor e_1 \rfloor^{il}_{il+l-1} + \Big(\sum\nolimits_{k=1}^{p-2} \lfloor e_{k+1,k} \rfloor^{il+k}_{il+l-1+k} \Big) \\
&\quad + \lfloor e_{p-1} \rfloor^{il+(p-1)}_{il+l-1+(p-1)} + \Big(\sum\nolimits_{k=1}^{p-2} \lfloor e_{p-k-1,p-k} \rfloor^{il+(p-1)+k}_{il+l-1+(p-1)+k} \Big) \Big) \Big) \\
&= \sum\nolimits_{i\geq 0} \Big(\lfloor e_1 \circ e_{1,1} \rfloor^{il}_{il+l} + \Big(\sum\nolimits_{k=1}^{p-2} \lfloor e_{k+1,k} \circ e_{k,k+1} \rfloor^{il+k}_{il+l+k} \Big) \\
&\quad + \lfloor e_{p-1} \circ e_{p-1,p-1} \rfloor^{il+(p-1)}_{il+l+(p-1)} + \Big(\sum\nolimits_{k=1}^{p-2} \lfloor e_{p-k-1,p-k} \circ e_{p-k,p-k-1} \rfloor^{il+(p-1)+k}_{il+l+(p-1)+k} \Big) \Big) \\
&\quad + \sum\nolimits_{i\geq 1} \lfloor e_{1,1} \circ e_1 \rfloor^{il-1}_{il+l-1} + \sum\nolimits_{i\geq 0} \Big(\Big(\sum\nolimits_{k=1}^{p-2} \lfloor e_{k,k+1} \circ e_{k+1,k} \rfloor^{il+k-1}_{il+l+k-1} \Big) \\
&\quad + \lfloor e_{p-1,p-1} \circ e_{p-1} \rfloor^{il-1+(p-1)}_{il+l-1+(p-1)} + \Big(\sum\nolimits_{k=1}^{p-2} \lfloor e_{p-k,p-k-1} \circ e_{p-k-1,p-k} \rfloor^{il-1+(p-1)+k}_{il+l-1+(p-1)+k} \Big) \Big) \\
&\overset{*}{=} \sum\nolimits_{i\geq 0} \Big(\lfloor e_{1,1} + e_{1,2} \circ e_{2,1} \rfloor^{il}_{il+l} + \Big(\sum\nolimits_{k=1}^{p-3} \lfloor e_{k+1,k} \circ e_{k,k+1} + e_{k+1,k+2} \circ e_{k+2,k+1} \rfloor^{il+k}_{il+l+k} \Big) \\
&\quad + \lfloor e_{p-1,p-2} \circ e_{p-2,p-1} + e_{p-1,p-1} \rfloor^{il+p-2}_{il+l+p-2} + \lfloor e_{p-1,p-1} + e_{p-1,p-2} \circ e_{p-2,p-1} \rfloor^{il+p-1}_{il+l+p-1} \\
&\quad + \Big(\sum\nolimits_{k=1}^{p-3} \lfloor e_{p-k-1,p-k} \circ e_{p-k,p-k-1} + e_{p-k-1,p-k-2} \circ e_{p-k-2,p-k-1} \rfloor^{il+p-1+k}_{il+l+p-1+k} \Big) \\
&\quad + \lfloor e_{1,2} \circ e_{2,1} + e_{1,1} \rfloor^{(i+1)l-1}_{(i+1)l+l-1} \Big) \overset{\text{L.}7(a)}{=} 0
\end{aligned}$$

In the step marked by "$*$" we sort the summands by their targets. Note that when splitting sums of the form $\sum_{k=1}^{p-2}(\ldots)_k$ into $(\ldots)_1 + \sum_{k=2}^{p-2}(\ldots)_k$ or into $(\ldots)_{p-2} + \sum_{k=1}^{p-3}(\ldots)_k$, the existence of the summand that is split off is ensured by $p \geq 3$.

(g): We first show that the differentials of the complex $\operatorname{Hom}^*(\operatorname{PRes}\mathbb{F}_p, \mathbb{F}_p)$ (cf. Lemma 19) are all zero: By Lemma 7, $\{\varepsilon\}$ is an $\mathbb{F}_p$-basis of $\operatorname{Hom}_{\mathbb{F}_p\mathrm{S}_p}(P_1, \mathbb{F}_p)$, and for $k \in [2, p-1]$ we have $\operatorname{Hom}_{\mathbb{F}_p\mathrm{S}_p}(P_k, \mathbb{F}_p) = 0$. So the only non-trivial $(d_k)^*$ are those where $\operatorname{Pr}_k = \operatorname{Pr}_{k+1} = P_1$. This is the case only when $k = lj + l - 1$ for some $j \geq 0$. Then $d_k = e_{1,1}$. For $\varepsilon \in \operatorname{Hom}(P_1, \mathbb{F}_p)$, we have $(d_k)^*(\varepsilon) = (-1)^k \varepsilon \circ e_{1,1} \overset{\text{L.}7(a)}{=} 0$. As $\operatorname{Hom}(P_1, \mathbb{F}_p) = \langle\varepsilon\rangle_{\mathbb{F}_p}$, we have $(d_k)^* = 0$.

So $\mathrm{H}^k \operatorname{Hom}^*(\operatorname{PRes}\mathbb{F}_p, \mathbb{F}_p) = \operatorname{Hom}^k(\operatorname{PRes}\mathbb{F}_p, \mathbb{F}_p)$. We use Lemma 19.

For $k = jl$, $j \geq 0$, we have $\bar{\Psi}^k(\overline{\iota^j}) \overset{(a)}{=} \varepsilon$, and $\{\varepsilon\}$ is a basis of $\mathrm{H}^k \operatorname{Hom}^*(\operatorname{PRes}\mathbb{F}_p, \mathbb{F}_p)$.

For $k = jl+l-1$, $j \geq 0$, we have $\bar{\Psi}^k(\overline{\chi\iota^j}) \overset{(e)}{=} \varepsilon$, and $\{\varepsilon\}$ is a basis of $\mathrm{H}^k \operatorname{Hom}^*(\operatorname{PRes}\mathbb{F}_p, \mathbb{F}_p)$. Finally, for $k = jl + r$ for some $j \geq 0$ and some $r \in [1, l-2]$ and for $k < 0$, we have $\mathrm{H}^k \operatorname{Hom}^*(\operatorname{PRes}\mathbb{F}_p, \mathbb{F}_p) = \{0\}$. □

2.3 An A_∞-structure on $\operatorname{Ext}^*_{\mathbb{F}_p\mathrm{S}_p}(\mathbb{F}_p, \mathbb{F}_p)$ as a minimal model of $\operatorname{Hom}^*_{\mathbb{F}_p\mathrm{S}_p}(\operatorname{PRes}\mathbb{F}_p, \operatorname{PRes}\mathbb{F}_p)$

Recall that $p \geq 3$ is a prime. Write $A := \operatorname{Hom}^*_{\mathbb{F}_p\mathrm{S}_p}(\operatorname{PRes}\mathbb{F}_p, \operatorname{PRes}\mathbb{F}_p)$, which becomes an A_∞-algebra $(A, (m_n)_{n\geq 1})$ over $R = \mathbb{F}_p$ via Lemma 13. We implement $\operatorname{Ext}^*_{\mathbb{F}_p\mathrm{S}_p}(\mathbb{F}_p, \mathbb{F}_p)$ as $\operatorname{Ext}^*_{\mathbb{F}_p\mathrm{S}_p}(\mathbb{F}_p, \mathbb{F}_p) := \mathrm{H}^* A$.

Our goal in this section is to construct an A_∞-structure $(m'_n)_{n\geq 1}$ on H^*A and a morphism of A_∞-algebras $f = (f_n)_{n\geq 1} : (\mathrm{H}^*A, (m'_n)_{n\geq 1}) \to (A, (m_n)_{n\geq 1})$ which satisfy the statements of Theorem 16. I.e. we will construct a minimal model of A. In preparation of the definitions of the f_n and m'_n, we name and examine certain elements of A:

Lemma 21. *Suppose given $k \in [2, p-1]$. We set*

$$\gamma_k := \sum\nolimits_{i\geq 0} \left(\lfloor e_k \rfloor^{k-1+li}_{k(l-1)+li} + \lfloor e_{p-k} \rfloor^{k-1+(p-1)+li}_{k(l-1)+(p-1)+li} \right) \in A^{k(l-2)+1}.$$

For $j \geq 0$, we have

$$\gamma_k \iota^j := \gamma_k \circ \iota^j = \iota^j \circ \gamma_k = \sum\nolimits_{i\geq 0} \left(\lfloor e_k \rfloor^{k-1+li}_{k(l-1)+l(i+j)} + \lfloor e_{p-k} \rfloor^{k-1+(p-1)+li}_{k(l-1)+(p-1)+l(i+j)} \right) \in A^{k(l-2)+1+jl}$$

and

$$\begin{aligned} m_1(\gamma_k \iota^j) = \sum\nolimits_{i\geq 0} \Big(& \lfloor e_{k-1,k} \rfloor^{k-2+li}_{k(l-1)+l(i+j)} + \lfloor e_{p-k+1,p-k} \rfloor^{k-2+(p-1)+li}_{k(l-1)+(p-1)+l(i+j)} \\ & + \lfloor e_{k,k-1} \rfloor^{k-1+li}_{k(l-1)+1+l(i+j)} + \lfloor e_{p-k,p-(k-1)} \rfloor^{k-1+(p-1)+li}_{k(l-1)+p+l(i+j)} \Big). \end{aligned}$$

Proof. We need to prove that γ_k is well-defined. Let $i \geq 0$.
We consider the first term. The complex $\operatorname{PRes}\mathbb{F}_p$ (cf. (3), (1)) has entry P_k at position $k(l-1)+li$ and at position $k-1+li$: We have $k(l-1)+li = (k-1+i)l + l - k$. So $\omega(k(l-1)+li) = l - (l-k) = k$ since $p-1 \leq l-k \leq l-1$. We have $\omega(k-1+li) = (k-1)+1 = k$ since $0 \leq k-1 \leq p-2$. As $k(l-1)+li, k-1+li \geq 0$, we have $\operatorname{Pr}_{k(l-1)+li} = P_{\omega(k(l-1)+li)} = P_k$ and $\operatorname{Pr}_{k-1+li} = P_{\omega(k-1+li)} = P_k$. So the first term is well-defined.
Now consider the second term. The complex $\operatorname{PRes}\mathbb{F}_p$ has entry P_{p-k} at position $k(l-1)+(p-1)+li$ and at position $k-1+(p-1)+li$: We have $k(l-1)+(p-1)+li = (i+k)l + (p-1) - k$, so $\omega(k(l-1)+(p-1)+li) = (p-1)-k+1 = p-k$ since $0 \leq (p-1)-k \leq p-2$. We have $\omega(k-1+(p-1)+li) = 2(p-1)-(k-1)-(p-1) = p-k$ since $p-1 \leq k-1+(p-1) \leq 2(p-1)-1$. As $k(l-1)+(p-1)+li, k-1+(p-1)+li \geq 0$, we have

$\mathrm{Pr}_{k(l-1)+(p-1)+li} = P_{\omega(k(l-1)+(p-1)+li)} = P_{p-k}$ and $\mathrm{Pr}_{k-1+(p-1)+li} = P_{\omega(k-1+(p-1)+li)} = P_{p-k}$. So the second term is well-defined.

The degree of the tuple of maps is computed to be $(k(l-1)+li)-(k-1+li) = k(l-2)+1 = (k(l-1)+(p-1)+li)-(k-1+(p-1)+li)$.

The explicit formula for $\gamma_k\iota^j$ is an application of Proposition 20(b).

The degree $|\gamma_k\iota^j| = k(l-2)+1$ is odd, so

$$\begin{aligned}
m_1(\gamma_k\iota^j) &\overset{\mathrm{L.13}}{=} d\circ\gamma_k\iota^j + \gamma_k\iota^j\circ d\\
&\overset{\mathrm{R.14}}{=} \sum\nolimits_{i\geq 0}\lfloor e_{\omega(k-2),\omega(k-1)}\rfloor^{k-2+li}_{k-1+li}\circ\sum\nolimits_{i\geq 0}\lfloor e_k\rfloor^{k-1+li}_{k(l-1)+l(i+j)}\\
&\quad+\sum\nolimits_{i\geq 0}\lfloor e_{\omega(p-1+k-2),\omega(p-1+k-1)}\rfloor^{k-2+(p-1)+li}_{k-1+(p-1)+li}\circ\sum\nolimits_{i\geq 0}\lfloor e_{p-k}\rfloor^{k-1+(p-1)+li}_{k(l-1)+(p-1)+l(i+j)}\\
&\quad+\sum\nolimits_{i\geq 0}\lfloor e_k\rfloor^{k-1+li}_{k(l-1)+l(i+j)}\circ\sum\nolimits_{i\geq 0}\lfloor e_{\omega(l-k),\omega(l-k+1)}\rfloor^{k(l-1)+l(i+j)}_{k(l-1)+1+l(i+j)}\\
&\quad+\sum\nolimits_{i\geq 0}\lfloor e_{p-k}\rfloor^{k-1+(p-1)+li}_{k(l-1)+(p-1)+l(i+j)}\circ\sum\nolimits_{i\geq 0}\lfloor e_{\omega(p-1-k),\omega(p-k)}\rfloor^{k(l-1)+(p-1)+l(i+j)}_{k(l-1)+p+l(i+j)}\\
&= \sum\nolimits_{i\geq 0}\lfloor e_{k-1,k}\rfloor^{k-2+li}_{k(l-1)+l(i+j)}+\sum\nolimits_{i\geq 0}\lfloor e_{p-k+1,p-k}\rfloor^{k-2+(p-1)+li}_{k(l-1)+(p-1)+l(i+j)}\\
&\quad+\sum\nolimits_{i\geq 0}\lfloor e_{k,k-1}\rfloor^{k-1+li}_{k(l-1)+1+l(i+j)}+\sum\nolimits_{i\geq 0}\lfloor e_{p-k,p-(k-1)}\rfloor^{k-1+(p-1)+li}_{k(l-1)+p+l(i+j)}
\end{aligned}$$

Note that in the second line $k-2+li\geq 0$ as $i\geq 0$ and $k\geq 2$. □

Lemma 22. *For* $j,j'\geq 0$*, we have* $\chi\iota^j\circ\chi\iota^{j'} = m_1(\gamma_2\iota^{j+j'})$.

Proof. It suffices to prove that $\chi\circ\chi = m_1(\gamma_2)$ since then $\chi\iota^j\circ\chi\iota^{j'} \overset{\mathrm{P.20}(e)}{=} \chi\circ\chi\circ\iota^{j+j'} = m_1(\gamma_2)\circ\iota^{j+j'} \overset{\mathrm{P.20}(c)}{=} m_1(\gamma_2\iota^{j+j'})$.
To determine when a composite is zero, we will need the following. For $0\leq k,k'<l$, we examine the condition

$$il+l-1+k = i'l+k'. \tag{8}$$

If $k=0$ then (8) holds iff $i=i'$ and $k'=l-1$.
If $k\geq 1$ then (8) holds iff $i+1=i'$ and $k'=k-1$.
So

$$\begin{aligned}
\chi\circ\chi \overset{p\geq 3}{=} &\Big(\sum\nolimits_{i\geq 0}\Big(\lfloor e_1\rfloor^{il}_{il+l-1}+\lfloor e_{2,1}\rfloor^{il+1}_{il+l}+\Big(\sum\nolimits_{k=2}^{p-2}\lfloor e_{k+1,k}\rfloor^{il+k}_{il+l-1+k}\Big)\\
&+\lfloor e_{p-1}\rfloor^{il+(p-1)}_{il+l-1+(p-1)}+\lfloor e_{p-2,p-1}\rfloor^{il+p}_{il+l+p-1}+\Big(\sum\nolimits_{k=2}^{p-2}\lfloor e_{p-k-1,p-k}\rfloor^{il+(p-1)+k}_{il+l-1+(p-1)+k}\Big)\Big)\Big)\\
&\circ\Big(\sum\nolimits_{i'\geq 0}\Big(\lfloor e_1\rfloor^{i'l}_{i'l+l-1}+\Big(\sum\nolimits_{k'=1}^{p-3}\lfloor e_{k'+1,k'}\rfloor^{i'l+k'}_{i'l+l-1+k'}\Big)+\lfloor e_{p-1,p-2}\rfloor^{i'l+p-2}_{i'l+l+p-3}\\
&+\lfloor e_{p-1}\rfloor^{i'l+(p-1)}_{i'l+l-1+(p-1)}+\Big(\sum\nolimits_{k'=1}^{p-3}\lfloor e_{p-k'-1,p-k'}\rfloor^{i'l+(p-1)+k'}_{i'l+l-1+(p-1)+k'}\Big)+\lfloor e_{1,2}\rfloor^{i'l+l-1}_{i'l+l+2(p-2)}\Big)\Big)
\end{aligned}$$

$$\begin{aligned}
&= \sum_{i\geq 0} \Big(\lfloor e_1 \circ e_{1,2} \rfloor^{il}_{il+l+2(p-2)} + \lfloor e_{2,1} \circ e_1 \rfloor^{il+1}_{il+2l-1} \\
&\quad + \Big(\sum_{k=2}^{p-2} \underbrace{\lfloor e_{k+1,k} \circ e_{k,k-1} \rfloor^{il+k}_{il+2l-1+k-1}}_{=0 \text{ by L.7(c)}} \Big) + \lfloor e_{p-1} \circ e_{p-1,p-2} \rfloor^{il+(p-1)}_{il+2l+p-3} \\
&\quad + \lfloor e_{p-2,p-1} \circ e_{p-1} \rfloor^{il+p}_{il+2l+p-2} + \Big(\sum_{k=2}^{p-2} \underbrace{\lfloor e_{p-k-1,p-k} \circ e_{p-k,p-k+1} \rfloor^{il+(p-1)+k}_{il+2l-1+p-1+k-1}}_{=0 \text{ by L.7(c)}} \Big) \Big) \\
&= \sum_{i\geq 0} \Big(\lfloor e_{1,2} \rfloor^{il}_{(i+2)l-2} + \lfloor e_{2,1} \rfloor^{il+1}_{(i+2)l-1} + \lfloor e_{p-1,p-2} \rfloor^{il+p-1}_{(i+2)l+p-3} + \lfloor e_{p-2,p-1} \rfloor^{il+p}_{(i+2)l+p-2} \Big) \\
&\overset{\text{L.21}}{=} m_1(\gamma_2)
\end{aligned}$$

□

Below are the definitions which will give a minimal A$_\infty$-algebra structure on H^*A and a quasi-isomorphism of A$_\infty$-algebras $\mathrm{H}^*A \to A$.

Definition 23. Recall from Proposition 20 that $\mathfrak{B} = \{\overline{\iota^j} \mid j \geq 0\} \sqcup \{\overline{\chi\iota^j} \mid j \geq 0\} = \{\overline{\chi^a\iota^j} \mid j \geq 0, a \in \{0,1\}\}$ is a basis of H^*A. For $n \in \mathbb{Z}_{\geq 1}$, we set

$$\mathfrak{B}^{\otimes n} := \{\overline{\chi^{a_1}\iota^{j_1}} \otimes \ldots \otimes \overline{\chi^{a_n}\iota^{j_n}} \in (\mathrm{H}^*A)^{\otimes n} \mid a_i \in \{0,1\} \text{ and } j_i \in \mathbb{Z}_{\geq 0} \text{ for all } i \in [1,n]\},$$

which is a basis of $(\mathrm{H}^*A)^{\otimes n}$ consisting of homogeneous elements.

For $n \geq 1$, we define the $\mathbb{F}_p$-linear map $f_n : (\mathrm{H}^*A)^{\otimes n} \to A$ as follows:

Case $n = 1$**:** f_1 is given on $\mathfrak{B}$ by $f_1(\overline{\iota^j}) := \iota^j$ and $f_1(\overline{\chi\iota^j}) := \chi\iota^j$.

Case $n \in [2, p-1]$**:** f_n is given on elements of $\mathfrak{B}^{\otimes n}$ by

$$f_n(\overline{\chi^{a_1}\iota^{j_1}} \otimes \ldots \otimes \overline{\chi^{a_n}\iota^{j_n}}) := \begin{cases} 0 & \text{if } \exists i \in [1,n] : a_i = 0 \\ (-1)^{n-1}\gamma_n\iota^{j_1+\ldots+j_n} & \text{if } 1 = a_1 = a_2 = \ldots = a_n \end{cases}$$

Case $n \geq p$**:** We set $f_n := 0$.

For $n \geq 1$, we define the $\mathbb{F}_p$-linear map $m'_n : (\mathrm{H}^*A)^{\otimes n} \to \mathrm{H}^*A$ by defining it on elements $\overline{\chi^{a_1}\iota^{j_1}} \otimes \ldots \otimes \overline{\chi^{a_n}\iota^{j_n}} \in \mathfrak{B}^{\otimes n}$:

Case $\exists i \in [1,n] : a_i = 0$**:**
$m'_n(\overline{\chi^{a_1}\iota^{j_1}} \otimes \ldots \otimes \overline{\chi^{a_n}\iota^{j_n}}) := 0$ for $n \neq 2$ and
$m'_2(\overline{\chi^{a_1}\iota^{j_1}} \otimes \overline{\chi^{a_2}\iota^{j_2}}) := \overline{\chi^{a_1+a_2}\iota^{j_1+j_2}}$ (Note that $a_1 + a_2 \in \{0,1\}$).

Case $a_1 = a_2 = \ldots = a_n = 1$**:**
$m'_n(\overline{\chi\iota^{j_1}} \otimes \ldots \otimes \overline{\chi\iota^{j_n}}) := 0$ for $n \neq p$ and
$m'_p(\overline{\chi\iota^{j_1}} \otimes \ldots \otimes \overline{\chi\iota^{j_p}}) := (-1)^p\overline{\iota^{p-1+j_1+\ldots+j_p}} = -\overline{\iota^{p-1+j_1+\ldots+j_p}}$.

Note that since $p \geq 3$, we have $m'_2(\overline{\chi\iota^{j_1}} \otimes \overline{\chi\iota^{j_2}}) = 0$ for $j_1, j_2 \geq 0$.

Theorem 24. *The pair* $(\mathrm{H}^*A,(m'_n)_{n\geq1})$ *is a minimal* A_∞*-algebra. The tuple* $(f_n)_{n\geq1}$ *is an quasi-isomorphism of* A_∞*-algebras from* $(\mathrm{H}^*A,(m'_n)_{n\geq1})$ *to* $(A,(m_n)_{n\geq1})$*. More precisely,* $f_1:(\mathrm{H}^*A,m'_1)\to(A,m_1)$ *induces the identity in homology.*

The proof of Theorem 24 will take the remainder of section 2.3. We will use Lemma 17.

Lemma 25. *The maps* f_n *and* m'_n *have degree* $|f_n|=1-n$ *and* $|m'_n|=2-n$*. I.e.* $(f_n)_{n\geq1}$ *is a pre-*A_∞*-morphism from* H^*A *to* A*, and* $(\mathrm{H}^*A,(m'_n)_{n\geq1})$ *is a pre-*A_∞*-algebra.*

Proof. We have $|f_1|=0$ as $|\overline{\iota^j}|=|\iota^j|$ and $|\overline{\chi\iota^j}|=|\chi\iota^j|$. For $n\geq p$ the map f_n is of degree $1-n$ as $f_n=0$. For $n\in[2,p-1]$ the statement $|f_n|=1-n$ is proven by checking the degrees for the elements of the basis $\mathfrak{B}^{\otimes n}$ whose image under f_n is non-zero:

$$\begin{aligned}|f_n(\overline{\chi\iota^{j_1}}\otimes\ldots\otimes\overline{\chi\iota^{j_n}})| &= |(-1)^{n-1}\gamma_n\iota^{j_1+\ldots+j_n}| \overset{\text{L.21}}{=} (j_1+\ldots+j_n)l+n(l-1)+1-n\\ &= 1-n+\sum\nolimits_{x=1}^{n}|\overline{\chi\iota^{j_x}}| = 1-n+|\overline{\chi\iota^{j_1}}\otimes\ldots\otimes\overline{\chi\iota^{j_n}}|\end{aligned}$$

Thus $|f_n|=1-n$ for all n and we have proven the first statement.

Now we show $|m'_n|=2-n$. As before, we only need check the degrees for basis elements whose image is non-zero: For $\overline{\chi^{a_1}\iota^{j_1}}\otimes\overline{\chi^{a_2}\iota^{j_2}}$, $j_1,j_2\geq0$, $a_1,a_2\in\{0,1\}$, $0\in\{a_1,a_2\}$, we have

$$\begin{aligned}|m'_2(\overline{\chi^{a_1}\iota^{j_1}}\otimes\overline{\chi^{a_2}\iota^{j_2}})| &= |\overline{\chi^{a_1+a_2}\iota^{j_1+j_2}}| = (a_1+a_2)(l-1)+l(j_1+j_2)\\ &= a_1(l-1)+j_1l+a_2(l-1)+j_2l = |\overline{\chi^{a_1}\iota^{j_1}}\otimes\overline{\chi^{a_2}\iota^{j_2}}|+(2-2).\end{aligned}$$

For $\overline{\chi\iota^{j_1}}\otimes\cdots\otimes\overline{\chi\iota^{j_p}}$, $j_x\geq0$ for $x\in[1,p]$, we have

$$\begin{aligned}|m'_p(\overline{\chi\iota^{j_1}}\otimes\cdots\otimes\overline{\chi\iota^{j_p}})| &= |\overline{\iota^{p-1+j_1+\ldots+j_p}}| = l(p-1+j_1+\ldots+j_p)\\ &= lp-l+l(j_1+\ldots+j_p) = lp-2p+2+l(j_1+\ldots+j_p)\\ &= p(l-1)+l(j_1+\ldots+j_p)+2-p = |\overline{\chi\iota^{j_1}}\otimes\cdots\otimes\overline{\chi\iota^{j_p}}|+2-p\end{aligned}$$

□

Lemma 26. *We have* $m'_1=0$*. The equation* (6)[1] *holds. The complex morphism* $f_1:(A',m'_1)\to(A,m_1)$ *is a quasi-isomorphism inducing the identity in homology.*

Proof. The equality $m'_1=0$ follows immediately from the definition. Thus $m_1\circ f_1=0=f_1\circ m'_1$. Moreover f_1 is a quasi-isomorphism inducing the identity in homology by construction, cf. Proposition 20(g). □

Lemma 27. *The map* f_1 *is injective.*

Proof. The set $X:=\{\chi^a\iota^j\mid a\in\{0,1\},j\in\mathbb{Z}_{\geq1}\}\subseteq A$ is linearly independent, since it consists of nonzero elements of different summands of the direct sum $A=\bigoplus_{k\in\mathbb{Z}}\mathrm{Hom}^k(\mathrm{PRes}\,\mathbb{F}_p,\mathrm{PRes}\,\mathbb{F}_p)$. The set $\mathfrak{B}$, which is a basis of H^*A, is mapped bijectively to X by f_1, so f_1 is injective. □

Lemma 28. *The equation* (6)[2] *holds.*

Proof. As $m'_1 = 0$, equation (6)[2] is equivalent to (cf. (7))

$$f_1 \circ m'_2 = m_1 \circ f_2 + m_2 \circ (f_1 \otimes f_1).$$

We check this equation on $\mathfrak{B}^{\otimes 2}$: Recall Proposition 20 and Definition 23.

$$\begin{aligned}
f_1 m'_2(\overline{\iota^j} \otimes \overline{\iota^{j'}}) &= \iota^{j+j'} = m_2(f_1 \otimes f_1)(\overline{\iota^j} \otimes \overline{\iota^{j'}}) = (m_1 \circ f_2 + m_2 \circ (f_1 \otimes f_1))(\overline{\iota^j} \otimes \overline{\iota^{j'}}) \\
f_1 m'_2(\overline{\iota^j} \otimes \overline{\chi\iota^{j'}}) &= \chi\iota^{j+j'} = m_2(f_1 \otimes f_1)(\overline{\iota^j} \otimes \overline{\chi\iota^{j'}}) \\
&= (m_1 \circ f_2 + m_2 \circ (f_1 \otimes f_1))(\overline{\iota^j} \otimes \overline{\chi\iota^{j'}}) \\
f_1 m'_2(\overline{\chi\iota^j} \otimes \overline{\iota^{j'}}) &= \chi\iota^{j+j'} = m_2(f_1 \otimes f_1)(\overline{\chi\iota^j} \otimes \overline{\iota^{j'}}) \\
&= (m_1 \circ f_2 + m_2 \circ (f_1 \otimes f_1))(\overline{\chi\iota^j} \otimes \overline{\iota^{j'}}) \\
f_1 m'_2(\overline{\chi\iota^j} \otimes \overline{\chi\iota^{j'}}) &= 0 \overset{\text{L.22}}{=} -m_1(\gamma_2 \iota^{j+j'}) + m_2(f_1 \otimes f_1)(\overline{\chi\iota^j} \otimes \overline{\chi\iota^{j'}}) \\
&= (m_1 \circ f_2 + m_2 \circ (f_1 \otimes f_1))(\overline{\chi\iota^j} \otimes \overline{\chi\iota^{j'}})
\end{aligned}$$

Note that there are no additional signs due to the Koszul sign rule since $|f_1| = 0$. ☐

The following results directly from Definition 23.

Corollary 29. *For* $n \geq 2$ *and* $a_1, \dots, a_n \in \{0,1\}$, $j_1, \dots, j_n \geq 0$, *we have*

$$f_n(\overline{\chi^{a_1}\iota^{j_1}} \otimes \dots \otimes \overline{\chi^{a_n}\iota^{j_n}}) = f_n(\overline{\chi^{a_1}} \otimes \dots \otimes \overline{\chi^{a_n}}) \circ \iota^{j_1 + \dots + j_n}.$$

If there is additionally an $x \in [1,n]$ *with* $a_x = 0$ *then*

$$f_n(\overline{\chi^{a_1}\iota^{j_1}} \otimes \dots \otimes \overline{\chi^{a_n}\iota^{j_n}}) = 0.$$

Equation (6)[n] can be reformulated as

$$\begin{aligned}
&\underbrace{f_1 \circ m'_n + \sum_{\substack{n=r+s+t \\ r,t \geq 0, s \geq 1 \\ s \leq n-1}} (-1)^{rs+t} f_{r+1+t} \circ (1^{\otimes r} \otimes m'_s \otimes 1^{\otimes t})}_{=:\Phi_n} \\
&= \underbrace{m_1 \circ f_n + \sum_{\substack{2 \leq r \leq n \\ i_1 + \dots + i_r = n \\ i_s \geq 1}} (-1)^v m_r \circ (f_{i_1} \otimes f_{i_2} \otimes \dots \otimes f_{i_r})}_{=:\Xi_n},
\end{aligned}$$

where $v = \sum_{1 \leq t < s \leq r} (1 - i_s) i_t$.

A term of the form $f_{r+1+t} \circ (1^{\otimes r} \otimes m'_s \otimes 1^{\otimes t})$, $s \geq 3$, $r+t \geq 1$, is zero because of Corollary 29 and the definition of m'_p. Also recall $m'_1 = 0$. Thus

$$\Phi_n = \sum_{\substack{n=r+2+t \\ r,t\geq 0}} (-1)^{2r+t} f_{n-1} \circ (1^{\otimes r} \otimes m'_2 \otimes 1^{\otimes t}) = \sum_{r=0}^{n-2} (-1)^{n-r} f_{n-1} \circ (1^{\otimes r} \otimes m'_2 \otimes 1^{\otimes n-r-2}). \tag{9}$$

Because of $m_k = 0$ for $k \geq 3$, we have

$$\Xi_n = \sum_{\substack{i_1+i_2=n \\ i_1,i_2\geq 1}} (-1)^{(1-i_2)i_1} m_2 \circ (f_{i_1} \otimes f_{i_2}) = \sum_{i=1}^{n-1} (-1)^{ni} m_2 \circ (f_i \otimes f_{n-i}). \tag{10}$$

We have proven:

Lemma 30. *For $n \geq 1$, condition* (6)[n] *is equivalent to $f_1 \circ m'_n + \Phi_n = m_1 \circ f_n + \Xi_n$ where Φ_n and Ξ_n are as in* (9) *and* (10)*.*

Lemma 31. *Condition* (6)[n] *holds for $n \geq 3$ and arguments $\overline{\chi^{a_1}\iota^{j_1}} \otimes \ldots \otimes \overline{\chi^{a_n}\iota^{j_n}} \in \mathfrak{B}^{\otimes n} = \{\overline{\chi^{a_1}\iota^{j_1}} \otimes \ldots \otimes \overline{\chi^{a_n}\iota^{j_n}} \in (\mathrm{H}^*A)^{\otimes n} \mid a_i \in \{0,1\}$ and $j_i \in \mathbb{Z}_{\geq 0}$ for all $i \in [1,n]\}$ where $0 \in \{a_1,\ldots,a_n\}$.*

Proof. Because of Lemma 30 and Definition 23 it is sufficient to show that

$$\Phi_n(\overline{\chi^{a_1}\iota^{j_1}} \otimes \ldots \otimes \overline{\chi^{a_n}\iota^{j_n}}) = \Xi_n(\overline{\chi^{a_1}\iota^{j_1}} \otimes \ldots \otimes \overline{\chi^{a_n}\iota^{j_n}})$$

if at least one a_x equals 0.

Case 1 At least two a_x equal 0:

To show $\Phi_n(\overline{\chi^{a_1}\iota^{j_1}} \otimes \ldots \otimes \overline{\chi^{a_n}\iota^{j_n}}) = 0$, we show $f_{n-1}(1^{\otimes r} \otimes m'_2 \otimes 1^{\otimes n-r-2})(\overline{\chi^{a_1}\iota^{j_1}} \otimes \ldots \otimes \overline{\chi^{a_n}\iota^{j_n}}) = 0$ for $r \in [0,n-2]$: In case both components of the argument of m'_2 are of the form $\overline{\chi^0\iota^j}$, the result of m'_2 is of the form $\overline{\iota^{j'}}$ (see Definition 23). Since $2 \leq n-1$, Corollary 29 implies the result of f_{n-1} is zero. Otherwise at least one of the components of the argument of f_{n-1} must be of the form $\overline{\iota^j}$ and the result of f_{n-1} is zero as well. So $\Phi_n(\overline{\chi^{a_1}\iota^{j_1}} \otimes \ldots \otimes \overline{\chi^{a_n}\iota^{j_n}}) = 0$. To show $\Xi_n(\overline{\chi^{a_1}\iota^{j_1}} \otimes \ldots \otimes \overline{\chi^{a_n}\iota^{j_n}}) = 0$, we show $m_2(f_i \otimes f_{n-i})(\overline{\chi^{a_1}\iota^{j_1}} \otimes \ldots \otimes \overline{\chi^{a_n}\iota^{j_n}}) = 0$ for $i \in [1,n-1]$:

- Suppose given $i \in [2,n-2]$: The statements $a_1 = \ldots = a_i = 1$ and $a_{i+1} = \ldots = a_n = 1$ cannot be true at the same time, so $f_i(\ldots) = 0$ or $f_{n-i}(\ldots) = 0$ and we have $m_2(f_i \otimes f_{n-i})(\overline{\chi^{a_1}\iota^{j_1}} \otimes \ldots \otimes \overline{\chi^{a_n}\iota^{j_n}}) = 0$.
- Suppose that $i = 1$. Because at least two a_x equal 0 the statement $a_2 = \ldots = a_n = 1$ cannot be true. Since $n-1 \geq 2$, we have $f_{n-1}(\ldots) = 0$ and $m_2(f_1 \otimes f_{n-1})(\overline{\chi^{a_1}\iota^{j_1}} \otimes \ldots \otimes \overline{\chi^{a_n}\iota^{j_n}}) = 0$.
- The case $i = n-1$ is analogous to the case $i = 1$.

So we have $\Phi_n(\overline{\chi^{a_1}\iota^{j_1}}\otimes\dots\otimes\overline{\chi^{a_n}\iota^{j_n}}) = 0 = \Xi_n(\overline{\chi^{a_1}\iota^{j_1}}\otimes\dots\otimes\overline{\chi^{a_n}\iota^{j_n}})$.

Case 2a Exactly one a_x equals 0, where $x\in[2,n-1]$.
We have $\Phi_n(\overline{\chi^{a_1}\iota^{j_1}}\otimes\dots\otimes\overline{\chi^{a_n}\iota^{j_n}}) = 0$: In case $n\geq p+1$, it follows from $f_{n-1}=0$. Let us check the case $n\in[3,p]$: Because of Definition 23, we have $f_{n-1}(1^{\otimes r}\otimes m_2'\otimes 1^{\otimes n-r-2})(\overline{\chi^{a_1}\iota^{j_1}}\otimes\dots\otimes\overline{\chi^{a_n}\iota^{j_n}}) = 0$ unless $r\in\{x-2,x-1\}$. So

$$
\begin{aligned}
&\Phi_n(\overline{\chi^{a_1}\iota^{j_1}}\otimes\dots\otimes\overline{\chi^{a_n}\iota^{j_n}})\\
&\quad= (-1)^{n-x+2} f_{n-1}(1^{\otimes x-2}\otimes m_2'\otimes 1^{\otimes n-x} - 1^{\otimes x-1}\otimes m_2'\otimes 1^{n-x-1})\\
&\qquad(\overline{\chi^{a_1}\iota^{j_1}}\otimes\dots\otimes\overline{\chi^{a_n}\iota^{j_n}})\\
&\quad= (-1)^{n-x} f_{n-1}(\overline{\chi\iota^{j_1}}\otimes\dots\otimes\overline{\chi\iota^{j_{x-2}}}\otimes m_2'(\overline{\chi\iota^{j_{x-1}}}\otimes\overline{\iota^{j_x}})\otimes\overline{\chi\iota^{j_{x+1}}}\otimes\dots\otimes\overline{\chi\iota^{j_n}}\\
&\qquad-\overline{\chi\iota^{j_1}}\otimes\dots\otimes\overline{\chi\iota^{j_{x-1}}}\otimes m_2'(\overline{\iota^{j_x}}\otimes\overline{\chi\iota^{j_{x+1}}})\otimes\overline{\chi\iota^{j_{x+2}}}\otimes\dots\otimes\overline{\chi\iota^{j_n}})\\
&\quad= (-1)^{n-x} f_{n-1}(\overline{\chi\iota^{j_1}}\otimes\dots\otimes\overline{\chi\iota^{j_{x-2}}}\otimes\overline{\chi\iota^{j_{x-1}+j_x}}\otimes\overline{\chi\iota^{j_{x+1}}}\otimes\dots\otimes\overline{\chi\iota^{j_n}}\\
&\qquad-\overline{\chi\iota^{j_1}}\otimes\dots\otimes\overline{\chi\iota^{j_{x-1}}}\otimes\overline{\chi\iota^{j_x+j_{x+1}}}\otimes\overline{\chi\iota^{j_{x+2}}}\otimes\dots\otimes\overline{\chi\iota^{j_n}})\\
&\quad= (-1)^{n-x}((-1)^{n-2}\gamma_{n-1}\iota^{j_1+\dots+j_n} - (-1)^{n-2}\gamma_{n-1}\iota^{j_1+\dots+j_n}) = 0
\end{aligned}
$$

To show $\Xi_n(\overline{\chi^{a_1}\iota^{j_1}}\otimes\dots\otimes\overline{\chi^{a_n}\iota^{j_n}}) = 0$, we show $m_2(f_i\otimes f_{n-i})(\overline{\chi^{a_1}\iota^{j_1}}\otimes\dots\otimes\overline{\chi^{a_n}\iota^{j_n}}) = 0$ for $i\in[1,n-1]$: The element $\chi^{a_x}\iota^{j_x}$ is a tensor factor of the argument of f_i or of f_{n-i}. We write $y=i$ or $y=n-i$ accordingly. Then $y\geq 2$ since $x\notin\{1,n\}$, so $f_y(\dots)=0$ and thus $m_2(f_i\otimes f_{n-i})(\overline{\chi^{a_1}\iota^{j_1}}\otimes\dots\otimes\overline{\chi^{a_n}\iota^{j_n}}) = 0$.
So $\Phi_n(\overline{\chi^{a_1}\iota^{j_1}}\otimes\dots\otimes\overline{\chi^{a_n}\iota^{j_n}}) = 0 = \Xi_n(\overline{\chi^{a_1}\iota^{j_1}}\otimes\dots\otimes\overline{\chi^{a_n}\iota^{j_n}})$.

Case 2b Only $a_1=0$, all other a_x equal 1.
We have $f_{n-1}(1^{\otimes r}\otimes m_2'\otimes 1^{\otimes n-r-2})(\overline{\chi^{a_1}\iota^{j_1}}\otimes\dots\otimes\overline{\chi^{a_n}\iota^{j_n}}) = 0$ unless $r=0$. So

$$
\begin{aligned}
\Phi_n(\overline{\chi^{a_1}\iota^{j_1}}\otimes\dots\otimes\overline{\chi^{a_n}\iota^{j_n}}) &= (-1)^n f_{n-1}(1^{\otimes 0}\otimes m_2'\otimes 1^{\otimes n-2})(\overline{\chi^{a_1}\iota^{j_1}}\otimes\dots\otimes\overline{\chi^{a_n}\iota^{j_n}})\\
&= (-1)^n f_{n-1}(m_2'(\overline{\iota^{j_1}}\otimes\overline{\chi\iota^{j_2}})\otimes\overline{\chi\iota^{j_3}}\otimes\dots\otimes\overline{\chi\iota^{j_n}})\\
&= (-1)^n f_{n-1}(\overline{\chi\iota^{j_1+j_2}}\otimes\overline{\chi\iota^{j_3}}\otimes\dots\otimes\overline{\chi\iota^{j_n}})\\
&= \begin{cases}\gamma_{n-1}\iota^{j_1+\dots+j_n} & 3\leq n\leq p\\ 0 & n\geq p+1\end{cases}
\end{aligned}
$$

We have $(f_i\otimes f_{n-1})(\overline{\chi^{a_1}\iota^{j_1}}\otimes\dots\otimes\overline{\chi^{a_n}\iota^{j_n}}) = 0$ if $i\geq 2$. So

$$
\begin{aligned}
\Xi_n(\overline{\chi^{a_1}\iota^{j_1}}\otimes\dots\otimes\overline{\chi^{a_n}\iota^{j_n}}) &= (-1)^{1\cdot n} m_2(f_1\otimes f_{n-1})(\overline{\chi^{a_1}\iota^{j_1}}\otimes\dots\otimes\overline{\chi^{a_n}\iota^{j_n}})\\
&\overset{(4)}{=} (-1)^n m_2\left((-1)^{n\cdot|\iota^{j_1}|} f_1(\overline{\iota^{j_1}})\otimes f_{n-1}(\overline{\chi\iota^{j_2}}\otimes\dots\otimes\overline{\chi\iota^{j_n}})\right)\\
&= (-1)^n m_2\left(\iota^{j_1}\otimes f_{n-1}(\overline{\chi\iota^{j_2}}\otimes\dots\otimes\overline{\chi\iota^{j_n}})\right)\\
&= \begin{cases}(-1)^n m_2(\iota^{j_1}\otimes(-1)^{n-2}\gamma_{n-1}\iota^{j_2+\dots+j_n}) & 3\leq n\leq p\\ 0 & n\geq p+1\end{cases}\\
&= \begin{cases}\gamma_{n-1}\iota^{j_1+\dots+j_n} & 3\leq n\leq p\\ 0 & n\geq p+1\end{cases}
\end{aligned}
$$

So $\Phi_n(\overline{\chi^{a_1}\iota^{j_1}}\otimes\ldots\otimes\overline{\chi^{a_n}\iota^{j_n}}) = \Xi_n(\overline{\chi^{a_1}\iota^{j_1}}\otimes\ldots\otimes\overline{\chi^{a_n}\iota^{j_n}})$.

Case 2c Only $a_n = 0$, all other a_x equal 1.
Argumentation analogous to case 2b gives

$$\begin{aligned}\Phi_n(\overline{\chi^{a_1}\iota^{j_1}}\otimes\ldots\otimes\overline{\chi^{a_n}\iota^{j_n}}) &= (-1)^2 f_{n-1}(1^{\otimes n-2}\otimes m_2'\otimes 1^{\otimes 0})(\overline{\chi^{a_1}\iota^{j_1}}\otimes\ldots\otimes\overline{\chi^{a_n}\iota^{j_n}})\\ &\overset{|m_2'|=0}{=} f_{n-1}(\overline{\chi\iota^{j_1}}\otimes\ldots\otimes\overline{\chi\iota^{j_{n-2}}}\otimes m_2'(\overline{\chi\iota^{j_{n-1}}}\otimes\overline{\iota^{j_n}}))\\ &= \begin{cases}(-1)^{n-2}\gamma_{n-1}\iota^{j_1+\ldots+j_n} & 3\le n\le p\\ 0 & n\ge p+1\end{cases}\end{aligned}$$

and

$$\begin{aligned}\Xi_n(\overline{\chi^{a_1}\iota^{j_1}}\otimes\ldots\otimes\overline{\chi^{a_n}\iota^{j_n}}) &= (-1)^{n(n-1)} m_2(f_{n-1}\otimes f_1)(\overline{\chi^{a_1}\iota^{j_1}}\otimes\ldots\otimes\overline{\chi^{a_n}\iota^{j_n}})\\ &\overset{|f_1|=0}{=} m_2\left(f_{n-1}(\overline{\chi\iota^{j_1}}\otimes\ldots\otimes\overline{\chi\iota^{j_{n-1}}})\otimes f_1(\overline{\iota^{j_n}})\right)\\ &= \begin{cases}(-1)^{n-2}\gamma_{n-1}\iota^{j_1+\ldots+j_n} & 3\le n\le p\\ 0 & n\ge p+1\end{cases}\end{aligned}$$

So $\Phi_n(\overline{\chi^{a_1}\iota^{j_1}}\otimes\ldots\otimes\overline{\chi^{a_n}\iota^{j_n}}) = \Xi_n(\overline{\chi^{a_1}\iota^{j_1}}\otimes\ldots\otimes\overline{\chi^{a_n}\iota^{j_n}})$.

□

Now we examine the cases where $a_1=\ldots=a_n=1$:

Lemma 32. *For* $n\ge 3$, *we have* $\Phi_n(\overline{\chi\iota^{j_1}}\otimes\ldots\otimes\overline{\chi\iota^{j_n}})=0$ *for* $\overline{\chi\iota^{j_1}}\otimes\ldots\otimes\overline{\chi\iota^{j_n}}\in\mathfrak{B}^{\otimes n}=\{\overline{\chi^{a_1}\iota^{j_1}}\otimes\ldots\otimes\overline{\chi^{a_n}\iota^{j_n}}\in(\mathrm{H}^*A)^{\otimes n}\mid a_i\in\{0,1\}$ *and* $j_i\in\mathbb{Z}_{\ge 0}$ *for all* $i\in[1,n]\}$.

Proof. We have $\Phi_n(\overline{\chi\iota^{j_1}}\otimes\ldots\otimes\overline{\chi\iota^{j_n}})=0$ since $\Phi_n=\sum_{r=0}^{n-2}(-1)^{n-r}f_{n-1}(1^{\otimes r}\otimes m_2'\otimes 1^{\otimes n-r-2})$ and the argument of m_2' is always of the form $\overline{\chi\iota^x}\otimes\overline{\chi\iota^y}$, whence its result is zero. □

Lemma 33. *Condition (6)[n] holds for* $n\in[3,p-1]$ *and arguments* $\overline{\chi\iota^{j_1}}\otimes\ldots\otimes\overline{\chi\iota^{j_n}}\in\mathfrak{B}^{\otimes n}=\{\overline{\chi^{a_1}\iota^{j_1}}\otimes\ldots\otimes\overline{\chi^{a_n}\iota^{j_n}}\in(\mathrm{H}^*A)^{\otimes n}\mid a_i\in\{0,1\}$ *and* $j_i\in\mathbb{Z}_{\ge 0}$ *for all* $i\in[1,n]\}$.

Proof. For computing Ξ_n, we first show that $m_2(f_k\otimes f_{n-k})(\overline{\chi\iota^{j_1}}\otimes\ldots\otimes\overline{\chi\iota^{j_n}})=0$ for $k\in[2,n-2]$. We will need the following congruence.

$$\begin{aligned}\underbrace{(k(l-1)+l(i+x))}_{\equiv_{p-1}k(l-1)+(p-1)+l(i+x)}-\underbrace{(n-k-1+li')}_{\equiv_{p-1}n-k-1+(p-1)+li'} &\equiv_{p-1} -k+k-n+1=-(n-1)\\ &\not\equiv_{p-1} 0 \qquad (11)\end{aligned}$$

The last statement results from $2\le n\le p-1$. We set "±" as a symbol for the (a posteriori irrelevant) signs in the following calculation. For $k\in[2,n-2]$, we have

$$m_2(f_k\otimes f_{n-k})(\overline{\chi\iota^{j_1}}\otimes\ldots\otimes\overline{\chi\iota^{j_n}})$$

$$
\begin{aligned}
&= \pm\, m_2((-1)^{k-1}\gamma_k\iota^{j_1+\ldots+j_k} \otimes (-1)^{n-k-1}\gamma_{n-k}\iota^{j_{k+1}+\ldots+j_n})\\
&\overset{\substack{j_1+\ldots+j_k=:x,\\ j_{k+1}+\ldots+j_n=:y}}{=} \pm\, \gamma_k\iota^x \circ \gamma_{n-k}\iota^y\\
&= \pm\Big(\sum\nolimits_{i\geq 0} \lfloor e_k\rfloor^{k-1+li}_{k(l-1)+l(i+x)} + \sum\nolimits_{i\geq 0}\lfloor e_{p-k}\rfloor^{k-1+(p-1)+li}_{k(l-1)+(p-1)+l(i+x)}\Big)\\
&\quad\circ\Big(\sum\nolimits_{i'\geq 0}\lfloor e_{n-k}\rfloor^{n-k-1+li'}_{(n-k)(l-1)+l(i'+y)} + \sum\nolimits_{i'\geq 0}\lfloor e_{p-n+k}\rfloor^{n-k-1+(p-1)+li'}_{(n-k)(l-1)+(p-1)+l(i'+y)}\Big)\\
&\overset{(11)}{=} 0.
\end{aligned}
$$

So

$$
\begin{aligned}
&\Xi_n(\overline{\chi\iota^{j_1}}\otimes\ldots\otimes\overline{\chi\iota^{j_n}})\\
&\qquad = m_2((-1)^n f_1\otimes f_{n-1} + (-1)^{n(n-1)} f_{n-1}\otimes f_1)(\overline{\chi\iota^{j_1}}\otimes\ldots\otimes\overline{\chi\iota^{j_n}})\\
&\qquad = m_2((-1)^{n+n|\overline{\chi\iota^{j_1}}|} f_1(\overline{\chi\iota^{j_1}})\otimes f_{n-1}(\overline{\chi\iota^{j_2}}\otimes\ldots\otimes\overline{\chi\iota^{j_n}})\\
&\qquad\quad + f_{n-1}(\overline{\chi\iota^{j_1}}\otimes\ldots\otimes\overline{\chi\iota^{j_{n-1}}})\otimes f_1(\overline{\chi\iota^{j_n}}))\\
&\qquad = m_2(\chi\iota^{j_1}\otimes(-1)^{n-2}\gamma_{n-1}\iota^{j_2+\ldots+j_n} + (-1)^{n-2}\gamma_{n-1}\iota^{j_1+\ldots+j_{n-1}}\otimes\chi\iota^{j_n})\\
&\qquad = (-1)^n(\chi\iota^{j_1}\circ\gamma_{n-1}\iota^{j_2+\ldots+j_n} + \gamma_{n-1}\iota^{j_1+\ldots+j_{n-1}}\circ\chi\iota^{j_n})\\
&\qquad \overset{\text{P.20(e),L.21}}{=} (-1)^n(\chi\circ\gamma_{n-1} + \gamma_{n-1}\circ\chi)\circ\iota^{j_1+\ldots+j_n}
\end{aligned}
$$

$$
\begin{aligned}
\chi\circ\gamma_{n-1} &= \Big(\sum\nolimits_{i\geq 0}\Big(\lfloor e_1\rfloor^{il}_{(i+1)l-1} + \Big(\sum\nolimits_{k=1}^{p-2}\lfloor e_{k+1,k}\rfloor^{il+k}_{(i+1)l-1+k}\Big)\\
&\quad + \lfloor e_{p-1}\rfloor^{il+(p-1)}_{(i+1)l-1+(p-1)} + \Big(\sum\nolimits_{k=1}^{p-2}\lfloor e_{p-k-1,p-k}\rfloor^{il+(p-1)+k}_{(i+1)l-1+(p-1)+k}\Big)\Big)\Big)\\
&\quad \circ\Big(\sum\nolimits_{i'\geq 0}\lfloor e_{n-1}\rfloor^{n-2+li'}_{(n-1)(l-1)+li'} + \sum\nolimits_{i'\geq 0}\lfloor e_{p-n+1}\rfloor^{n-2+(p-1)+li'}_{(n-1)(l-1)+(p-1)+li'}\Big)\\
&\underset{\substack{k\rightsquigarrow n-1\\ i'\rightsquigarrow i+1}}{\overset{3\leq n\leq p-1}{=}} \sum\nolimits_{i\geq 0}\lfloor e_{n,n-1}\circ e_{n-1}\rfloor^{il+n-1}_{(n-1)(l-1)+l(i+1)}\\
&\quad + \sum\nolimits_{i\geq 0}\lfloor e_{p-n,p-n+1}\circ e_{p-n+1}\rfloor^{il+p-1+n-1}_{(n-1)(l-1)+(p-1)+l(i+1)}\\
&= \sum\nolimits_{i\geq 0}\Big(\lfloor e_{n,n-1}\rfloor^{il+n-1}_{n(l-1)+1+li} + \lfloor e_{p-n,p-n+1}\rfloor^{il+p-1+n-1}_{n(l-1)+p+li}\Big)\\
\gamma_{n-1}\circ\chi &= \Big(\sum\nolimits_{i'\geq 0}\lfloor e_{n-1}\rfloor^{n-2+li'}_{(n-1+i'-1)l+2(p-1)-(n-1)} + \sum\nolimits_{i'\geq 0}\lfloor e_{p-n+1}\rfloor^{n-2+(p-1)+li'}_{(n-1+i')l-n+p}\Big)\\
&\quad \circ\Big(\sum\nolimits_{i\geq 0}\Big(\lfloor e_1\rfloor^{il}_{(i+1)l-1} + \Big(\sum\nolimits_{k=1}^{p-2}\lfloor e_{k+1,k}\rfloor^{il+k}_{(i+1)l-1+k}\Big)\\
&\quad + \lfloor e_{p-1}\rfloor^{il+(p-1)}_{(i+1)l-1+(p-1)} + \Big(\sum\nolimits_{k=1}^{p-2}\lfloor e_{p-k-1,p-k}\rfloor^{il+(p-1)+k}_{(i+1)l-1+(p-1)+k}\Big)\Big)\Big)\\
&\overset{k\rightsquigarrow p-n}{=} \sum\nolimits_{i'\geq 0}\lfloor e_{n-1}\circ e_{n-1,n}\rfloor^{n-2+li'}_{(n-1+i')l-1+(p-1)+(p-n)}\\
&\quad + \sum\nolimits_{i'\geq 0}\lfloor e_{p-n+1}\circ e_{p-n+1,p-n}\rfloor^{n-2+(p-1)+li'}_{(n+i')l-1+p-n}\\
&= \sum\nolimits_{i'\geq 0}\lfloor e_{n-1,n}\rfloor^{n-2+li'}_{n(l-1)+i'l} + \sum\nolimits_{i'\geq 0}\lfloor e_{p-n+1,p-n}\rfloor^{n-2+(p-1)+li'}_{n(l-1)+(p-1)+i'l}
\end{aligned}
$$

So $\chi\circ\gamma_{n-1}+\gamma_{n-1}\circ\chi = m_1(\gamma_n)$ by Lemma 21. Therefore

$$\begin{aligned}\Xi_n(\overline{\chi\iota^{j_1}}\otimes\ldots\otimes\overline{\chi\iota^{j_n}}) &= (-1)^n m_1(\gamma_n)\circ\iota^{j_1+\ldots+j_n} \overset{\text{P.20(c)}}{=} (-1)^n m_1(\gamma_n\iota^{j_1+\ldots+j_n})\\ &= -\,m_1((-1)^{n-1}\gamma_n\iota^{j_1+\ldots+j_n})\\ &= -\,m_1\circ f_n(\overline{\chi\iota^{j_1}}\otimes\ldots\otimes\overline{\chi\iota^{j_n}}).\end{aligned}$$

We conclude using Lemma 30 by

$$(f_1\circ m'_n+\Phi_n)(\overline{\chi\iota^{j_1}}\otimes\ldots\otimes\overline{\chi\iota^{j_n}}) \overset{\text{L.32,D.23}}{=} 0 = (m_1\circ f_n+\Xi_n)(\overline{\chi\iota^{j_1}}\otimes\ldots\otimes\overline{\chi\iota^{j_n}}).$$

□

Lemma 34. *Condition* (6)[p] *holds for arguments* $\overline{\chi\iota^{j_1}}\otimes\ldots\otimes\overline{\chi\iota^{j_p}}\in\mathfrak{B}^{\otimes p}=\{\overline{\chi^{a_1}\iota^{j_1}}\otimes\ldots\otimes\overline{\chi^{a_p}\iota^{j_p}}\in(\mathrm{H}^*A)^{\otimes p}\mid a_i\in\{0,1\}$ *and* $j_i\in\mathbb{Z}_{\geq 0}$ *for all* $i\in[1,p]\}$.

Proof. Recall that $|\iota| = l = 2(p-1)$ is even, $|\chi| = l-1$ is odd and $|f_i| = 1-i$ by Lemma 25. We have

$$\begin{aligned}\Xi_p(\overline{\chi\iota^{j_1}}\otimes\ldots\otimes\overline{\chi\iota^{j_p}}) &= \textstyle\sum_{i=1}^{p-1}(-1)^{pi}m_2(f_i\otimes f_{p-i})(\overline{\chi\iota^{j_1}}\otimes\ldots\otimes\overline{\chi\iota^{j_p}})\\ &= \textstyle\sum_{i=1}^{p-1}(-1)^{pi+i(1-(p-i))}m_2(f_i(\overline{\chi\iota^{j_1}}\otimes\ldots\otimes\overline{\chi\iota^{j_i}})\otimes f_{p-i}(\overline{\chi\iota^{j_{i+1}}}\otimes\ldots\otimes\overline{\chi\iota^{j_p}}))\\ &= \textstyle\sum_{i=1}^{p-1} f_i(\overline{\chi\iota^{j_1}}\otimes\ldots\otimes\overline{\chi\iota^{j_i}})\circ f_{p-i}(\overline{\chi\iota^{j_{i+1}}}\otimes\ldots\otimes\overline{\chi\iota^{j_p}})\\ &\overset{p\geq 3}{=} \chi\iota^{j_1}\circ(-1)^{p-2}\gamma_{p-1}\iota^{j_2+\ldots+j_p}+(-1)^{p-2}\gamma_{p-1}\iota^{j_1+\ldots+j_{p-1}}\circ\chi\iota^{j_p}\\ &\qquad+\textstyle\sum_{i=2}^{p-2}(-1)^{i-1}\gamma_i\iota^{j_1+\ldots+j_i}\circ(-1)^{p-i-1}\gamma_{p-i}\iota^{j_{i+1}+\ldots+j_p}\\ &\overset{\text{P.20(b)}}{=} (-1)^p\Big(\chi\circ\gamma_{p-1}+\gamma_{p-1}\circ\chi+\textstyle\sum_{k=2}^{p-2}\gamma_k\circ\gamma_{p-k}\Big)\circ\iota^{j_1+\ldots+j_p}\end{aligned}$$

$$\begin{aligned}\chi\circ\gamma_{p-1} &= \Big(\textstyle\sum_{i\geq 0}\Big(\lfloor e_1\rfloor^{il}_{(i+1)l-1}+\Big(\textstyle\sum_{k=1}^{p-2}\lfloor e_{k+1,k}\rfloor^{il+k}_{(i+1)l-1+k}\Big)\\ &\qquad+\lfloor e_{p-1}\rfloor^{il+(p-1)}_{(i+1)l-1+(p-1)}+\Big(\textstyle\sum_{k=1}^{p-2}\lfloor e_{p-k-1,p-k}\rfloor^{il+(p-1)+k}_{(i+1)l-1+(p-1)+k}\Big)\Big)\Big)\\ &\qquad\circ\Big(\textstyle\sum_{i'\geq 0}\lfloor e_{p-1}\rfloor^{(p-1)-1+li'}_{(p-1)(l-1)+li'}+\textstyle\sum_{i'\geq 0}\lfloor e_1\rfloor^{-1+2(p-1)+li'}_{(p-1)(l-1)+(p-1)+li'}\Big)\\ &= \textstyle\sum_{i\geq 0}\lfloor e_{p-1}\rfloor^{il+(p-1)}_{(p-1)(l-1)+l(i+1)}+\textstyle\sum_{i\geq 0}\lfloor e_1\rfloor^{il}_{(p-1)(l-1)+(p-1)+li}\\ &= \textstyle\sum_{i\geq 0}\lfloor e_{p-1}\rfloor^{il+(p-1)}_{(p+i-1)l+(p-1)}+\textstyle\sum_{i\geq 0}\lfloor e_1\rfloor^{il}_{(p+i-1)l}\end{aligned}$$

$$\begin{aligned}\gamma_{p-1}\circ\chi &= \Big(\textstyle\sum_{i'\geq 0}\lfloor e_{p-1}\rfloor^{(p-1)-1+li'}_{(p+i'-2)l+(p-1)}+\textstyle\sum_{i'\geq 0}\lfloor e_1\rfloor^{-1+2(p-1)+li'}_{(p+i'-1)l}\Big)\\ &\qquad\circ\Big(\textstyle\sum_{i\geq 0}\Big(\lfloor e_1\rfloor^{il}_{(i+1)l-1}+\Big(\textstyle\sum_{k=1}^{p-2}\lfloor e_{k+1,k}\rfloor^{il+k}_{(i+1)l-1+k}\Big)\end{aligned}$$

$$
\begin{aligned}
&+\lfloor e_{p-1}\rfloor^{il+(p-1)}_{(i+1)l-1+(p-1)} + \Big(\sum\nolimits_{k=1}^{p-2}\lfloor e_{p-k-1,p-k}\rfloor^{il+(p-1)+k}_{(i+1)l-1+(p-1)+k}\Big)\Big)\Big)\\
&= \sum\nolimits_{i'\geq 0}\lfloor e_{p-1}\rfloor^{(p-1)-1+li'}_{(p+i'-1)l-1+(p-1)} + \sum\nolimits_{i'\geq 0}\lfloor e_1\rfloor^{-1+2(p-1)+li'}_{(p+i')l-1}\\
&= \sum\nolimits_{i'\geq 0}\lfloor e_{p-1}\rfloor^{p-2+i'l}_{(p+i'-1)l+p-2} + \sum\nolimits_{i'\geq 0}\lfloor e_1\rfloor^{i'l+l-1}_{(p+i'-1)l+l-1}
\end{aligned}
$$

$$
\begin{aligned}
\gamma_k\circ\gamma_{p-k} &= \Big(\sum\nolimits_{i\geq 0}\lfloor e_k\rfloor^{k-1+li}_{(i+k-1)l+l-k} + \sum\nolimits_{i\geq 0}\lfloor e_{p-k}\rfloor^{k-1+(p-1)+li}_{(i+k)l+(p-1)-k}\Big)\\
&\quad\circ\Big(\sum\nolimits_{i'\geq 0}\lfloor e_{p-k}\rfloor^{p-k-1+li'}_{(p-k)(l-1)+li'} + \sum\nolimits_{i'\geq 0}\lfloor e_k\rfloor^{-k+2(p-1)+li'}_{(p-k)(l-1)+(p-1)+li'}\Big)\\
&= \sum\nolimits_{i\geq 0}\lfloor e_k\rfloor^{k-1+li}_{(p-k)(l-1)+(p-1)+l(i+k-1)} + \sum\nolimits_{i\geq 0}\lfloor e_{p-k}\rfloor^{k-1+(p-1)+li}_{(p-k)(l-1)+l(i+k)}\\
&= \sum\nolimits_{i\geq 0}\lfloor e_k\rfloor^{k-1+li}_{(p-k+i+k-1)l-(p-k)+(p-1)} + \sum\nolimits_{i\geq 0}\lfloor e_{p-k}\rfloor^{k-1+(p-1)+li}_{(p-k+i+k)l-(p-k)}\\
&= \sum\nolimits_{i\geq 0}\lfloor e_k\rfloor^{k-1+li}_{(p+i-1)l+k-1} + \sum\nolimits_{i\geq 0}\lfloor e_{p-k}\rfloor^{k-1+(p-1)+li}_{(p+i-1)l+k-1+(p-1)}\,.
\end{aligned}
$$

Thus

$$
\begin{aligned}
\chi\circ\gamma_{p-1}&+\gamma_{p-1}\circ\chi + \sum\nolimits_{k=2}^{p-2}\gamma_k\circ\gamma_{p-k}\\
&= \sum\nolimits_{i\geq 0}\sum\nolimits_{k=0}^{p-2}\Big(\lfloor e_{k+1}\rfloor^{k+li}_{(p+i-1)l+k} + \lfloor e_{p-k-1}\rfloor^{k+(p-1)+li}_{(p+i-1)l+k+(p-1)}\Big)\\
&= \sum\nolimits_{i\geq 0}\sum\nolimits_{k'=0}^{l-1}\lfloor e_{\omega(k')}\rfloor^{k'+li}_{(p-1+i)l+k'} \overset{\text{P.}20(a)}{=} \iota^{p-1}
\end{aligned}
$$

and

$$
\Xi_p(\overline{\chi\iota^{j_1}}\otimes\ldots\otimes\overline{\chi\iota^{j_p}}) = (-1)^p\iota^{p-1+j_1+\ldots+j_p}\,.
$$

So we conclude using Lemma 30 by

$$
\begin{aligned}
(f_1\circ m'_p+\Phi_p)(\overline{\chi\iota^{j_1}}\otimes\ldots\otimes\overline{\chi\iota^{j_p}}) &\overset{\text{L.}32,\text{D.}23}{=} (-1)^p\iota^{p-1+j_1+\ldots+j_p}\\
&\overset{\text{D.}23}{=} (m_1\circ f_p+\Xi_p)(\overline{\chi\iota^{j_1}}\otimes\ldots\otimes\overline{\chi\iota^{j_p}}).
\end{aligned}
$$

☐

Lemma 35. *Condition* (6)[n] *holds for* $n\in[p+1,2(p-1)]$ *and arguments* $\overline{\chi\iota^{j_1}}\otimes\ldots\otimes\overline{\chi\iota^{j_n}}\in\mathfrak{B}^{\otimes n}=\{\overline{\chi^{a_1}\iota^{j_1}}\otimes\ldots\otimes\overline{\chi^{a_n}\iota^{j_n}}\in(\mathrm{H}^*A)^{\otimes n}\mid a_i\in\{0,1\}$ *and* $j_i\in\mathbb{Z}_{\geq 0}$ *for all* $i\in[1,n]\}$.

Proof. As $f_k=0$ for $k\geq p$, we have

$$
\Xi_n(\overline{\chi\iota^{j_1}}\otimes\ldots\otimes\overline{\chi\iota^{j_n}}) = \sum\nolimits_{k=n-p+1}^{p-1}(-1)^{nk}m_2(f_k\otimes f_{n-k})(\overline{\chi\iota^{j_1}}\otimes\ldots\otimes\overline{\chi\iota^{j_n}})
$$

The right side is a linear combination of terms of the form $\gamma_k\circ\gamma_{n-k}$ for $k\in[n-p-1,p-1]$. We have

$$
\gamma_k\circ\gamma_{n-k} = \Big(\sum\nolimits_{i\geq 0}\lfloor e_k\rfloor^{k-1+li}_{k(l-1)+li} + \sum\nolimits_{i\geq 0}\lfloor e_{p-k}\rfloor^{k-1+(p-1)+li}_{k(l-1)+(p-1)+li}\Big)
$$

$$\circ \left(\sum\nolimits_{i'\geq 0} \lfloor e_{n-k} \rfloor^{n-k-1+li'}_{(n-k)(l-1)+li'} + \sum\nolimits_{i'\geq 0} \lfloor e_{p-n+k} \rfloor^{n-k-1+(p-1)+li'}_{(n-k)(l-1)+(p-1)+li'}\right)$$

A necessary condition for that term to be non-zero is $k(l-1) \equiv_{p-1} n-k-1$ as $l = 2(p-1)$. We have

$$k(l-1) - (n-k-1) \equiv_{p-1} -k - n + k + 1 = 1 - n \not\equiv_{p-1} 0,$$

since $p \leq n-1 \leq 2(p-1)-1$. So $\gamma_k \circ \gamma_{n-k} = 0$ and $\Xi_n(\overline{\chi\iota^{j_1}} \otimes \ldots \otimes \overline{\chi\iota^{j_n}}) = 0$. We conclude using Lemma 30 by

$$(f_1 \circ m'_n + \Phi_n)(\overline{\chi\iota^{j_1}} \otimes \ldots \otimes \overline{\chi\iota^{j_n}}) \overset{\text{L.32,D.23}}{=} 0 \overset{\text{D.23}}{=} (m_1 \circ f_n + \Xi_n)(\overline{\chi\iota^{j_1}} \otimes \ldots \otimes \overline{\chi\iota^{j_n}}).$$

□

One could formulate a lemma similar to Lemma 35 for the case $n > 2(p-1)$ as then the sum $\sum_{k=n-p+1}^{p-1} (-1)^{nk} m_2(f_k \otimes f_{n-k})(\overline{\chi\iota^{j_1}} \otimes \ldots \otimes \overline{\chi\iota^{j_n}})$ is in fact empty. Instead we use Lemma 18 to prove (6)[n] for $n > 2p-2$:

Proof of Theorem 24. Lemmas 26, 28, 31 and 33 to 35 ensure that (6)[n] holds for $n \in [1, 2p-2]$. Then Lemma 18 with $k = p$ proves that (6)[n] holds for all $n \in [1, \infty]$, cf. also Definition 23. By Lemma 27, f_1 is injective. By Lemma 25, the degrees are as required in Lemma 17. Lemma 17 proves that $(\mathrm{H}^* A, (m'_n)_{n\geq 1})$ is an A_∞-algebra and $(f_n)_{n\geq 1}$ is an A_∞-morphism from $(\mathrm{H}^* A, (m'_n)_{n\geq 1})$ to $(A, (m_n)_{n\geq 1})$. By Lemma 26, we have $m'_1 = 0$. Thus $(\mathrm{H}^* A, (m'_n)_{n\geq 1})$ is a minimal A_∞-algebra. By Lemma 26, the complex morphism $f_1 : (\mathrm{H}^* A, m'_1) \to (A, m_1)$ is a quasi-isomorphism which induces the identity in homology. So the A_∞-morphism $(f_n)_{n\geq 1} : (\mathrm{H}^* A, (m'_n)_{n\geq 1}) \to (A, (m_n)_{n\geq 1})$ is a quasi-isomorphism and the proof of Theorem 24 is complete. □

2.4 At the prime 2

We examine the case at the prime 2. We use a direct approach. Note that S_2 is a cyclic group so the theory of cyclic groups applies as well.

We have $\mathbb{F}_2\,\mathrm{S}_2 = \{0, (\mathrm{id}), (1,2), (\mathrm{id}) + (1,2)\}$. We have maps given by

$$\begin{array}{rrcl}
\varepsilon : & \mathbb{F}_2\,\mathrm{S}_2 & \longrightarrow & \mathbb{F}_2 \\
& a(\mathrm{id}) + b(1,2) & \longmapsto & a + b \\
D : & \mathbb{F}_2\,\mathrm{S}_2 & \longrightarrow & \mathbb{F}_2\,\mathrm{S}_2 \\
& a(\mathrm{id}) + b(1,2) & \longmapsto & (a+b)\left((\mathrm{id}) + (1,2)\right).
\end{array}$$

We see that ε is surjective and $\ker \varepsilon = \ker D = \operatorname{im} D = \{0, (\mathrm{id}) + (1,2)\}$. The maps ε and D are $\mathbb{F}_2\,\mathrm{S}_2$-linear, where $\mathbb{F}_2$ is the $\mathbb{F}_2\,\mathrm{S}_2$-module that corresponds to the trivial representation of S_2. So we have a projective resolution of $\mathbb{F}_2$ with augmentation ε by

$$\operatorname{PRes} \mathbb{F}_2 := (\cdots \xrightarrow{D} \underbrace{\mathbb{F}_2\,\mathrm{S}_2}_{1} \xrightarrow{D} \underbrace{\mathbb{F}_2\,\mathrm{S}_2}_{0} \to \underbrace{0}_{-1} \to \cdots),$$

where the degrees are written below.

We set e_1 to be the identity on $\mathbb{F}_2\,\mathrm{S}_2$.

Let $A := \mathrm{Hom}^*_{\mathbb{F}_2\,\mathrm{S}_2}(\mathrm{PRes}\,\mathbb{F}_2, \mathrm{PRes}\,\mathbb{F}_2)$ and let the A_∞-structure on A be $(m_n)_{n\geq 1}$ (cf. Lemma 13). Recall the conventions concerning $\mathrm{Hom}^k_B(C, C')$ for complexes C, C' and $k \in \mathbb{Z}$.

Lemma 36. *An $\mathbb{F}_2$-basis of H^*A is given by $\{\overline{\xi^j} \mid j \geq 0\}$ where*

$$\xi := \sum\nolimits_{i\geq 0} \lfloor e_1 \rfloor^i_{i+1} \in A.$$

Proof. Straightforward induction yields, for $j \geq 0$,

$$\xi^j = \sum\nolimits_{i\geq 0} \lfloor e_1 \rfloor^i_{i+j}\,.$$

We have

$$\begin{aligned} m_1(\xi^j) &= d \circ \xi^j - (-1)^j \xi^j \circ d = d \circ \xi^j + \xi^j \circ d \\ &= \Big(\sum\nolimits_{i\geq 0} \lfloor D \rfloor^i_{i+1}\Big) \circ \Big(\sum\nolimits_{i\geq 0} \lfloor e_1 \rfloor^i_{i+j}\Big) + \Big(\sum\nolimits_{i\geq 0} \lfloor e_1 \rfloor^i_{i+j}\Big) \circ \Big(\sum\nolimits_{i\geq 0} \lfloor D \rfloor^i_{i+1}\Big) \\ &= \sum\nolimits_{i\geq 0} \lfloor D \rfloor^i_{i+j+1} + \sum\nolimits_{i\geq 0} \lfloor D \rfloor^i_{i+j+1} = 0, \end{aligned}$$

so ξ^j is a cycle. As $\mathrm{Hom}_{\mathbb{F}_2\,\mathrm{S}_2}(\mathbb{F}_2\,\mathrm{S}_2, \mathbb{F}_2) = \{0, \varepsilon\}$ and $\varepsilon \circ D = 0$, the differentials of $\mathrm{Hom}^*(\mathrm{PRes}\,\mathbb{F}_2, \mathbb{F}_2)$ (cf. Lemma 19) are all zero. So $\{\varepsilon\}$ is an $\mathbb{F}_2$-basis of $\mathrm{H}^k\,\mathrm{Hom}^*(\mathrm{PRes}\,\mathbb{F}_2, \mathbb{F}_2)$ for $k \geq 0$. Since in the notion of Lemma 19, $\bar{\Psi}_k(\overline{\xi^k}) = \varepsilon$, the set $\{\overline{\xi^k}\}$ is an $\mathbb{F}_2$-basis of $\mathrm{H}^k\,\mathrm{Hom}^*(\mathrm{PRes}\,\mathbb{F}_2, \mathrm{PRes}\,\mathbb{F}_2)$ for $k \geq 0$. For $k < 0$ we have $\mathrm{H}^k\,\mathrm{Hom}^*(\mathrm{PRes}\,\mathbb{F}_2, \mathrm{PRes}\,\mathbb{F}_2) \cong \mathrm{H}^k\,\mathrm{Hom}^*(\mathrm{PRes}\,\mathbb{F}_2, \mathbb{F}_2) = 0$. So $\{\overline{\xi^j} \mid j \geq 0\}$ is an $\mathbb{F}_2$-basis of H^*A. □

We define families of maps $(f_n : (\mathrm{H}^*A)^{\otimes n} \to A)_{n\geq 1}$ and $(m'_n : (\mathrm{H}^*A)^{\otimes n} \to \mathrm{H}^*A)_{n\geq 1}$ as follows. f_1 and m'_2 are given on a basis by

$$\begin{aligned} f_1(\overline{\xi^j}) &:= \xi^j && \text{for } j \geq 0 \\ m'_2(\overline{\xi^j} \otimes \overline{\xi^k}) &:= \overline{\xi^{j+k}} && \text{for } j, k \geq 0. \end{aligned}$$

All other maps are set to zero.

It is straightforward to check that $(\mathrm{H}^*A, (m'_n)_{n\geq 1})$ is a pre-A_∞-algebra and $(f_n)_{n\geq 1}$ is a pre-A_∞-morphism from H^*A to A. As m'_2 is associative, $(\mathrm{H}^*A, (m'_n)_{n\geq 1})$ is a dg-algebra, so in particular an A_∞-algebra. As $f_k = 0$ for $k \neq 1$, (6)[n] simplifies to

$$f_1 \circ m'_n = m_n \circ (\underbrace{f_1 \otimes \cdots \otimes f_1}_{n \text{ factors}}).$$

As $m'_n = 0$ and $m_n = 0$ for $n \geq 3$, (6)[n] is satisfied for all $n \geq 3$. For $n \in \{1, 2\}$, we have

$$f_1 \circ m'_1 = m_1 \circ f_1$$

$$f_1 \circ m'_2 = m_2(f_1 \otimes f_1).$$

The second equation follows immediately from the definition of m'_2 and f_1. The first equation holds as $m'_1 = 0$ and the images of f_1 are all cycles. So (6)[n] holds for all n and $(f_n)_{n\geq 1}$ is an A_∞-morphism from $(\mathrm{H}^*A, (m'_n)_{n\geq 1})$ to $(A, (m_n)_{n\geq 1})$. By the construction of f_1, it induces the identity on homology. Thus $(\mathrm{H}^*A, (m'_n)_{n\geq 1})$ is a minimal model of $(A, (m_n)_{n\geq 1})$.

Remark 37 (Comparison with primes $p \geq 3$)**.** At a prime $p \geq 3$, we have constructed a projective resolution with period length $l = 2(p-1)$ in (2). If one constructs a projective resolution of $\mathbb{Z}_{(2)}$ analogous to the case $p \geq 3$, we have a sequence of the form

$$\cdots \to \mathbb{Z}_{(2)}\,\mathrm{S}_2 \xrightarrow{\hat{e}^*_{2,2}} \mathbb{Z}_{(2)}\,\mathrm{S}_2 \xrightarrow{\hat{e}_{2,2}} \mathbb{Z}_{(2)}\,\mathrm{S}_2 \xrightarrow{\hat{e}^*_{2,2}} \mathbb{Z}_{(2)}\,\mathrm{S}_2 \xrightarrow{\hat{e}_{2,2}} \mathbb{Z}_{(2)}\,\mathrm{S}_2 \to 0 \to \cdots$$

with a period length of 2, where

$$\begin{aligned} \hat{e}_{2,2}\colon\ (\mathrm{id}) &\longmapsto (\mathrm{id}) - (1,2) \\ \hat{e}^*_{2,2}\colon\ (\mathrm{id}) &\longmapsto (\mathrm{id}) + (1,2). \end{aligned}$$

However, modulo 2 the differentials $\hat{e}_{2,2}$ and $\hat{e}^*_{2,2}$ reduce to the same map $D : \mathbb{F}_2\,\mathrm{S}_2 \to \mathbb{F}_2\,\mathrm{S}_2$, so we obtain a period length of 1.

The maps ι resp. χ from Proposition 20 may be identified with ξ^2 resp. ξ. This way, the definition of m'_2 at the prime 2 is readily compatible with Definition 23.

References

[1] Benson, D.J. *Representations and cohomology I: Basic representation theory of finite groups and associative algebras*, Cambridge Univ. Press, 1998

[2] Bourbaki, N. *Éléments de mathématique - Algèbre - Chapitre 10: Algèbre homologique*, 1980

[3] Green, J.A. *Walking around the Brauer tree*, Journal of the Australian Mathematical Society, Volume 17, Issue 2, p. 197-213, 1974

[4] Gugenheim, V.K.A.M., Lambe, L.A., and Stasheff, J.D., *Perturbation theory in differential homological algebra II*, Illinois J. Math. 35, p. 357-373, 1991

[5] James, G.D., *The Representation Theory of the Symmetric Groups*, Lecture Notes in Mathematics 682, Springer-Verlag, 1978

[6] Johansson, L., and Lambe, L. *Transferring Algebra Structures Up to Homology Equivalence*, Math. Scand. 89, p. 181-200, 2001

[7] Kadeishvili, T.V., *On the homology theory of fiber spaces*, Russian Math. Surveys, 35:3, p. 231-238, 1980

[8] Kadeishvili, T.V., *Algebraic Structure in the Homologies of an $A(\infty)$-Algebra (Russian)*, Bulletin of the Academy of Sciences of the Georgian SSR 108 No 2, p. 249-252, 1982

[9] Kadeishvili, T.V., *The functor D for a category of $A(\infty)$-algebras (Russian)*, Bulletin of the Academy of Sciences of the Georgian SSR 125, p. 273-276, 1987

[10] Keller, B., *Introduction to A-infinity algebras and modules*, Homology, Homotopy and Applications vol. 3(1), 2001

[11] Keller, B., *Addendum to 'Introduction to A-infinity algebras and modules'*, 2002

[12] Klamt, A., A_∞*-structures on the algebra of extension of Verma modules in the parabolic category $\mathcal{O}$*, Diplomarbeit, 2010

[13] Künzer, M., *Ties for the integral group ring of the symmetric group*, Thesis, 1999

[14] Lefèvre-Hasegawa, K., *Sur les* A_∞*-catégories*, Thesis, 2003

[15] Madsen, D.,*Homological aspects in representation theory*, Thesis, 2002

[16] Merkulov, S.A., *Strong homotopy algebras of a Kähler manifold*, Int. Math. Res. Notices, Vol. 1999, p. 153-164, 1999

[17] Peel, M.H., *Hook representations of the symmetric groups*, Glasgow Mathematical Journal, 1971

[18] Prouté, A., *Algèbres différentielles fortement homotopiquement associatives*, Thèse d'Etat, Université Paris VII, 1984

[19] Roggenkamp, K. W., *Integral representations and structure of finite group rings*, Séminaire de mathématiques supérieures 71., Presses de l'Université de Montréal, 1980

[20] Stasheff, J.D., *Homotopy associativity of H-spaces II*, Trans. Amer. Math. Soc. 108, p. 293-312

[21] Vejdemo-Johansson, M., *Computation of* A_∞*-algebras in group cohomology*, Thesis, 2008

[22] Vejdemo-Johansson, M., *A partial* A_∞*-structure on the cohomology of* $C_m \times C_n$, Journal of Homotopy and Related Structures, vol. 3(1), p. 1-11, 2008

[23] Vejdemo-Johansson, M., *Blackbox computation of* A_∞*-algebras*, Georgian Mathematical Journal Volume 17 Issue 2, p. 391-404, 2010